

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 47/48	A1	(11) International Publication Number: WO 99/08711 (43) International Publication Date: 25 February 1999 (25.02.99)
(21) International Application Number: PCT/US98/16970 (22) International Filing Date: 13 August 1998 (13.08.98) (30) Priority Data: 60/054,743 13 August 1997 (13.08.97) US 09/132,808 12 August 1998 (12.08.98) US (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, R440, Emeryville, CA 94608 (US). (72) Inventors: ZUCKERMANN, Ronald, N.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). HUANG, Carol; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). MURPHY, John, E.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). UNO, Tetsuo; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). (74) Agents: FUJITA, Sharon, M. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94622-8097 (US).		(81) Designated States: JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: LIPID-POLYAMIDE CONJUGATES AND COMPOSITIONS FOR NUCLEIC ACID DELIVERY (57) Abstract In accordance with the present invention, there are provided lipid-conjugated polyamide compounds and related compositions and methods thereof. Lipid-conjugated polyamide compounds of the present invention are particularly useful as vehicles for delivering biologically active agents to a target site. In particular, the invention compounds are effective at facilitating the delivery of polynucleotides to cells. The present invention also provides a method for producing stable formulations of polynucleotides complexed with a delivery vehicle.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

LIPID-POLYAMIDE CONJUGATES AND COMPOSITIONS FOR NUCLEIC ACID DELIVERY

Field of the Invention

5 The present invention relates to lipid-conjugated polyamide compounds, methods for making them, as well as compositions and methods for their use, such as, for example, in the delivery of biologically active agents to cells.


Background of the Invention

10 The discovery of new therapeutic agents having increasingly complex molecular structure has presented new challenges related to how they can be efficiently delivered to targeted sites. For example, recent developments in recombinant DNA technology and human genome characterization have enabled identification of the molecular origins of many genetic and acquired diseases and construction of appropriate plasmids containing
 15 desired genes. However, the efficient delivery of these large and heavily charged constructs, having molecular weights of up to tens of millions of daltons and containing several tens of thousands of negative charges into cells remains a substantial challenge. Studies evaluating the use of neutral and cationic liposome structures as vehicles for the delivery of polynucleotides to cells have met with limited success, as these encapsulated
 20 structures are rather large and unstable.

 Accordingly, compounds that can be used as effective vehicles for the efficient delivery of large complex agents, such as polynucleotides, to cells would be highly desirable.

Summary of the Invention

25 The present invention is directed to lipid-conjugated polyamide compounds and compositions thereof that are particularly useful in the delivery of bioactive agents to cells. Specifically, the present invention provides lipid-conjugated polyamide compounds having the general formula:



wherein n is an integer selected from 1 to about 48 and m is an integer selected from about 2 to about 48,

wherein R_1 for each monomeric unit, $\text{---}(\overset{\overset{R_1}{|}}{\text{N}}\text{---}\overset{\overset{\text{O}}{\parallel}}{\text{C}})\text{---}$, and R_a are
 5 independently selected from the group consisting of a hydrogen atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group; -SH; an optionally substituted, branched or straight chain aliphatic group having from about 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and
 10 phosphorus, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or
 15 more double or triple bonds; and a lipid moiety that is optionally bonded to a linker moiety,

wherein R_1 is not a hydrogen atom for at least one monomeric unit,

wherein R_c is selected from a hydrogen atom; a hydroxy group; an amino group; a hydrazine group; a sulfonyl group; -SH; an optionally substituted, branched or straight
 20 chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted
 25 arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl group of said arylalkyl optionally has one or more double or triple bonds; and a lipid moiety that is optionally bonded to a linker moiety,

wherein when R_1 , R_a , or R_c is an aryl or arylalkyl group having fewer than about
 30 5 carbon atoms in a backbone structure, said backbone structure further comprises one or more heteroatoms,

wherein W for each monomeric unit is independently selected from an optionally substituted, branched or straight chain divalent moiety having from 1 to about 50 atoms and optionally, one or more double or triple bonds in a backbone that contains carbon and optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said optional substitution of W may be a lipid moiety that is optionally bonded to a linker moiety,

wherein said lipid moiety is a hydrophobic or amphipathic moiety selected from the group consisting of:

(i) optionally substituted aryl or arylalkyl moieties having from about 14 to about 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and

(ii) optionally substituted, branched or straight chain aliphatic moieties having from about 10 to about 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic moieties optionally have one or more double or triple bonds, and

wherein at least one of R_a , R_c , W for a single monomeric unit and R_1 for a single monomeric unit comprises a lipid moiety optionally bonded to a linker moiety.


The present invention also provides a method of synthesizing lipid-conjugated polyamide compounds, said method comprising:

a) contacting

(1) a lipid reactant, with

(2) an oligomer reactant, wherein said oligomer reactant has the

general formula:



wherein n is an integer selected from 1 to about 48, and m is an integer from about 2 to about 48,

wherein each T_a and T_c is independently selected from a terminal group and a reactive moiety that is capable of further reaction with said lipid reactant,

wherein R_1 for each monomeric unit, $\begin{array}{c} R_1 \\ | \\ N - W - C \\ || \\ O \end{array}$, in said oligomer reactant is selected from the group consisting of a hydrogen

atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group, -SH; an optionally substituted, branched or straight chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the
5 aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that
10 optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl group of said arylalkyl optionally has one or more double or triple bonds; and a reactive moiety that is capable of further reaction with said lipid reactant,

wherein when R_1 , R_a , or R_c is an aryl or arylalkyl group having
15 fewer than about 5 carbon atoms in a backbone structure, said backbone structure further comprises one or more heteroatoms,

wherein R_1 is not a hydrogen atom for at least one monomeric unit,

wherein W for each monomeric unit is selected from an optionally
20 substituted, branched or straight chain divalent moiety having from 1 to about 50 atoms in a backbone that contains carbon, and optionally contains nitrogen, oxygen, sulfur, and phosphorus, and optionally one or more double or triple bonds, wherein said optional substitution of W may be a reactive moiety that is capable of further reaction with said lipid
25 reactant,

wherein at least one of T_a , T_c , W for a single monomeric unit, or R_1 for a single monomeric unit comprises a reactive moiety that is capable of further reaction with said lipid reactant; then

b) reacting said lipid reactant with said oligomer reactant to conjugate the
30 lipid reactant to the oligomer reactant.

In another embodiment, the present invention provides a composition comprising a lipid-conjugated polyamide compound of the present invention and a biologically active agent.

In yet another embodiment, the present invention provides a method for inducing
5 the uptake of a biologically active agent by a cell, said method comprising:

providing a composition comprising an effective amount of a biologically active agent and a lipid-conjugated polyamide compound of the present invention; then
contacting a biological sample with an effective dose of said composition,
wherein said biological sample comprises a cell.

10 In still another embodiment, the present invention provides a method for inducing the uptake of a biologically active agent by a cell *in vivo*, said method comprising:

providing a composition comprising an effective amount of a biologically active agent and a lipid-conjugated polyamide compound of the present invention; then
administering an effective dose of said composition to a subject.

15 In a further embodiment, the present invention provides a method of expressing a gene in a mammal, said method comprising:

administering a polynucleic acid complexed with a lipid-conjugated polyamide compound of the present invention to a mammal,

20 wherein said polynucleic acid is capable of functionally expressing said gene in said mammal, and

wherein said complex is effective at transfecting said gene into a cell in said mammal.

In another embodiment, the present invention provides a method for substantially inhibiting nuclease-induced polynucleotide degradation, said method comprising:

25 contacting a polynucleotide with a degradation-inhibiting quantity of a lipid-conjugated polyamide compound, and

introducing the polynucleotide and the lipid-conjugated polyamide compound into a nuclease-containing environment.

In still a further embodiment, the present invention provides a method of making
30 a stable preparation of a polynucleic acid complexed with a delivery vehicle, said method comprising:

- a) providing a polynucleic acid in a first liquid carrier as a dilute polynucleic acid solution that is substantially precipitant-free;
- b) providing a delivery vehicle-forming compound in a second liquid carrier as a delivery vehicle solution that is substantially precipitant-free;
- 5 c) combining said dilute polynucleic acid solution with said delivery vehicle solution to form a dilute preparation of delivery vehicle/polynucleic acid complex; then
- d) reducing the volume of said dilute preparation to form a stable preparation of delivery vehicle/polynucleic acid complex,
- 10 wherein the concentration of polynucleic acid in said stable preparation is higher than the concentration of polynucleic acid in said dilute polynucleic acid solution, and
- wherein said stable preparation is substantially precipitant-free.

15 Brief Description of the Drawings

Figure 1 shows a reaction scheme for preparing lipid-conjugated polyamide compounds.

Figure 2 is a plasmid map of vector CMVkm2.

Figure 3 shows the effect of the +/- charge ratio of a complex of plasmid DNA (pCMVkmLUC) and a lipid-conjugated polyamide of the present invention (i.e.,
20 Compound 16 from Table 2) on the transfection of HT1080 cells. Luciferase activity in transfected cells is shown on the y-axis in (pg/20 μ l) and the +/- charge ratio is shown on the x-axis. The open bars refer to cells grown in FCS-supplemented medium and the shaded bars refer to cells grown in OptiMEM (i.e., serum-free medium).

Figure 4 shows the relationship between transfection efficiency and total number of monomeric units in the oligomeric moiety of lipid-conjugated polyamide compounds of the present invention. Luciferase activity (all normalized to luciferase activity corresponding to the 36-mer (i.e., n=3, m=12 with reference to formula (I))) in
25 transfected HT1080 cells is shown on the y-axis and length of oligomer is shown on the x-axis. The lipid-conjugated polyamide/DNA complexes had a +/- charge ratio of 2:1.
30

Figure 5 shows a comparison in transfection efficiency between DMPE-conjugated polyamide compounds (i.e., Compounds 16 and 23) and DOPE-conjugated polyamide compounds (i.e., Compounds 20 and 24). The open bars refer to cells grown in FCS-supplemented medium and the shaded bars refer to cells cultured in optiMEM (i.e., serum-free medium).

Figure 6 shows luciferase expression in Balb/C mouse lung, liver, and spleen tissue after *in vivo* transfection with lipid-conjugated polyamide compound (Compound 16)/pCMVkmLUC complex.

Figure 7 shows zeta potential stability over a period of 8 days of a formulation of lipid-conjugated polyamide (Compound 16)/DNA complex, prepared by the "dilution-concentration" formulation method of the present invention.

Figure 8 shows zeta potential stability over a period of 8 days of a formulation of lipid-conjugated polyamide (Compound 23)/DNA complex, prepared by the "dilution-concentration" formulation method.

Figure 9 shows particle size stability over a period of 8 days of a formulation of lipid-conjugated polyamide (Compound 16)/DNA complex, prepared by the "dilution-concentration" formulation method.

Figure 10 shows particle size stability over a period of 8 days of a formulation of lipid-conjugated polyamide (Compound 23)/DNA complex, prepared by the "dilution-concentration" formulation method.

Figure 11 illustrates the stability of a formulation of lipid-conjugated polyamide (Compound 16)/DNA complex prepared by the "dilution-concentration" formulation method and a formulation of DMRIE-CTM/DNA complex prepared by a conventional formulation method. HT1080 cells were transfected with lipid-conjugated polyamide/DNA complex 2 days postformulation and with DMRIE-CTM/DNA complex immediately after formulation. Results are shown for transfected cells cultured in both FCS-supplemented and OptiMEM media.

Figure 12 illustrates the stability of a formulation of lipid-conjugated polyamide (Compound 16)/DNA complex prepared by the "dilution-concentration" formulation method and a formulation of DMRIE-CTM/DNA complex prepared by a conventional formulation method. HT1080 cells were transfected with lipid-conjugated

polyamide/DNA complex 18 days postformulation and with DMRIE-C™/DNA complex immediately after formulation. Results are shown for transfected cells cultured in both FCS-supplemented and OptiMEM media.

Figure 13 illustrates the stability of a formulation of lipid-conjugated polyamide (Compound 23)/DNA complex prepared by the "dilution-concentration" formulation method and a formulation of DMRIE-C™/DNA complex prepared by a conventional formulation method. HT1080 cells were transfected with lipid-conjugated polyamide/DNA complex 4 days postformulation and with DMRIE-C™/DNA complex immediately after formulation. Results are shown for transfected cells cultured in both FCS-supplemented and OptiMEM media.


Figure 14 illustrates the stability of a formulation of lipid-conjugated polyamide (Compound 23)/DNA complex prepared by the "dilution-concentration" formulation method and a formulation of DMRIE-C™/DNA complex prepared by a conventional formulation method. HT1080 cells were transfected with lipid-conjugated polyamide/DNA complex 12 days postformulation and with DMRIE-C™/DNA complex immediately after formulation. Also shown are the effects of doubling the quantity of transfection medium on transfection efficiency. Results are shown for transfected cells cultured in both FCS-supplemented and OptiMEM media.

Figure 15 illustrates transfection efficiency using conventional and "dilution-concentration" formulation methods. "Mixing" refers to a conventional formulation method (in which the delivery vehicle and DNA were mixed immediately prior to transfection). "Preformed" refers to formulation of delivery vehicle/DNA complexes by the "dilution-concentration" formulation method, followed by transfection 1 to 5 days later. Also shown is cell toxicity. Results are shown for transfected cells cultured in FCS-supplemented medium.


General Methods and Detailed Description

The terms "lipid-conjugated polyamide compound" and "lipid-conjugated compound" are used interchangeably herein to refer to a compounds of the present invention which have both an oligomeric amide moiety and one or more lipid moieties.

The terms "oligomeric" and "oligomeric amide" are used interchangeably herein to refer to two or more monomer units that are linked together by an amide bond,



- 5 The term "monomer" or "monomeric" unit refers to the unit defined by the formula



As used herein, the term "lipid" refers to a hydrophobic or amphipathic moiety.

- 10 A lipid moiety can be conjugated directly to the oligomeric amide moiety, or optionally, indirectly to the oligomeric amide moiety via a linker moiety.

The terms "oligomeric reactant," "oligomer reactant," and "oligomeric amide reactant," and "lipid reactant" refer herein to reactive species from which lipid-conjugated polyamide compounds of the present invention are synthesized.

- 15 The term "backbone" refers herein to the scaffold structure of a moiety which is typically either a straight chain, branched or cyclic arrangement of covalently bonded carbon or heteroatoms (i.e., the scaffold structure does not include any of the hydrogen atoms, or alternatively, substitution groups bonded to it).

- As used herein, the term "optionally substituted" refers to the replacement of
20 hydrogen with a monovalent radical, such as, for example, hydroxyl-, carboxyl-, phospho-, amino-, halo-, alkyl-, aryl-, arylalkyl-, thioamido-, amido-, nitro-, cyano-, haloalkyl-, and the like.

- The term "aliphatic" refers herein to straight chain, branched and cyclic compounds that do not have aromatic properties, which contain carbon atoms and
25 optionally, one or more heteroatoms (i.e., one or more functional groups, such as, for example, a substituted amino, an alkoxy, a carbonyl, an ester, and the like), and optionally one or more double or triple bond (i.e., alkenyl or alkynyl, respectively).

- As used herein, the term "aryl" refers to aromatic groups, such as, for example, monocyclic and polycyclic aromatic groups, having one or more heteroatoms
30 incorporated therein (e.g., nitrogen, oxygen, sulfur, and phosphorus). The term "polycyclic" refers herein to both fused and non-fused cyclic structure in which at least

one cyclic structure is aromatic. Exemplary aryl moieties include, phenyl, naphthyl, and the like.

The term "arylalkyl" refers herein to an alkyl group, having optional functional groups incorporated therein, substituted with an aryl group. Exemplary arylalkyl
5 moieties include benzyl, picolyl, and the like.

As used herein, the term "delivery vehicle" refers to a compound and/or structure that complexes with and facilitates the delivery of a biologically active compound to a target site. Suitable delivery vehicles employed in the practice of the present invention include, for example, lipid-conjugated polyamide compounds of the present invention,
10 lipids, polycationic non-lipid compounds, liposomes, and the like.

As used herein, the term "complex" refers to a structure formed by interaction between two or more compounds or structures. Such interaction can be via chemical interaction, such as, for example, covalent, ionic, or secondary bonding (e.g., hydrogen bonding), and the like, or via physical interaction, such as, for example, encapsulation,
15 entrapment, and the like.


For example, lipid-conjugated polyamide compounds of the present invention can be complexed to a low molecular weight biologically active compound (including for example, oligonucleotides) via covalent bonding through an intermediately positioned sequence of amino acids that is susceptible to degradation by endogenous proteolytic
20 enzymes. Thus, for example, exposure of the complex to degradative enzymes results in cleavage and subsequent release of the biologically active compound from the complex. Lipid-conjugated polyamide compounds of the present invention can also be complexed to biologically active compounds, such as polynucleotides, via ionic or secondary bonding, or alternatively via encapsulation or entrapment.

25 The terms "polynucleotide" and "polynucleic acid" are used interchangeably herein to refer to DNA, RNA, and analogues thereof, peptide-nucleic acids, as well as DNA or RNA having non-phosphate containing nucleotides. Polynucleotides employed in the practice of the present invention can be single-stranded, double-stranded, or chimeric single- or double-stranded molecules.

All publications mentioned herein are incorporated herein by reference for the purpose of disclosing and describing features of the invention for which the publications are cited in connection with.

5 LIPID-CONJUGATED POLYAMIDE COMPOUNDS

The present invention provides lipid-conjugated polyamide compounds having the general formula:



10 wherein n is an integer selected from 1 to about 48 and m is an integer selected from about 2 to about 48,

wherein R_1 for each monomeric unit, $\left(\begin{array}{c} R_1 \\ | \\ N - W - C \\ || \\ O \end{array} \right)_n$, and R_a are independently selected from the group consisting of a hydrogen atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group; -SH; an optionally substituted, branched or
 15 straight chain aliphatic group having 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted
 20 arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and a lipid moiety that is optionally bonded to linker moiety,

wherein R_1 is not a hydrogen atom for at least one monomeric unit,

25 wherein R_c is selected from a hydrogen atom; a hydroxy group; an amino group; a hydrazine group; a sulfonyl group; -SH; an optionally substituted, branched or straight chain aliphatic group having 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl
 30 group having from about 3 to about 12 carbon atoms in a backbone structure that

optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl group of said arylalkyl optionally has one or more double or triple bonds; and a lipid moiety that is optionally bonded to a linker moiety,

wherein when R_1 , R_a , or R_c is an aryl or arylalkyl group having fewer than about 5 carbon atoms in a backbone structure, said backbone structure further comprises one or more heteroatoms, such as, for example, oxygen and/or nitrogen,

wherein W for each monomeric unit is independently selected from an optionally substituted, branched or straight chain divalent moiety having from 1 to about 50 atoms and optionally, one or more double or triple bonds in a backbone that contains carbon and optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said optional substitution of W may be a lipid moiety that is optionally bonded to a linker moiety,

wherein said lipid moiety is a hydrophobic or amphipathic moiety selected from the group consisting of:

(i) optionally substituted aryl or arylalkyl moieties having from about 14 to about 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl group of said arylalkyl optionally has one or more double or triple bonds; and

(ii) optionally substituted, branched or straight chain aliphatic moieties having from about 10 to about 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic moieties optionally have one or more double or triple bonds, and

wherein at least one of R_a , R_c , W for a single monomeric unit, and R_1 for a single monomeric unit comprises a lipid moiety.

Lipid-conjugated polyamides of the present invention can be random polymers where each R_1 and W varies from monomer to monomer (e.g., where n is 1 and m is an integer from about 2 to about 48). Alternatively, the lipid-conjugated polyamides can be polymers having m number of n-mers (i.e., where n is greater than 1 and m is an integer from about 2 to about 48) that are either repeating (i.e., each n-mer is the same) or

randomly variable (i.e., the monomer composition of each n-mer varies from n-mer to n-mer).

Typically, the integer n is not more than about 40, more typically not more than about 20, and even more typically not more than about 6. Preferably, n is about 3. The integer m is typically not more than about 40, more typically not more than about 25. Usually, the integer m is not more than about 15, typically not more than about 12, and even more typically not more than about 8.

Heteroatoms (i.e., nitrogen, oxygen, sulfur, and phosphorus) incorporated into the backbone structure of aliphatic moieties and the alkyl component of arylalkyl moieties typically form a functional group, such as, for example, a substituted amine, a carbonyl, an alkoxy, an ester, and the like. Thus, aliphatic and arylalkyl moieties employed in compounds of the present invention optionally have one or more functional groups incorporated therein. Heteroatoms incorporated into the backbone structure of aryl moieties are incorporated as ring atoms in cyclic aryl moieties.

When R_1 , R_a , and R_c are aliphatic, they typically contain at least 2 carbon atoms in a backbone structure and more typically contain at least about 3 carbon atoms in a backbone structure. Aryl and arylalkyl R_1 , R_a , and R_c groups can be linear or cyclic. Aryl and arylalkyl R_1 , R_a , and R_c having less than about 5 carbon atoms in a backbone structure, also typically have one or more heteroatoms in the backbone structure, such as, for example, nitrogen and/or oxygen. Typically aryl and arylalkyl R_1 , R_a , and R_c have at least about 5 carbon atoms in a backbone structure.

R_a is typically -OH, -H, -SH, -COOH, sulfonyl, or a lipid moiety optionally conjugated to a linker moiety. R_c is typically -OH, -H, -SH, -NH₂, sulfonyl, hydrazine, or a lipid moiety optionally conjugated to a linker moiety. Preferably, either R_a or R_c is a lipid moiety optionally conjugated to a linker moiety.

R_1 can be a sidechain that is cationic, anionic, or neutral at physiological relevant pH. Typically, physiological pH is at least about 5.5 and typically at least about 6.0. More typically, physiological pH is at least about 6.5. Usually, physiological pH is less than about 8.5 and typically less than about 8.0. More typically, physiological pH is less than about 7.5.

Suitable cationic sidechains include, for example, aminoalkyl (e.g., aminoethyl, aminopropyl, aminobutyl, aminopentyl, and the like) as well as derivatives thereof; (S)- α -methylethylenediamino and derivatives thereof; trimethylaminoethyl and derivatives thereof; guanidinoalkyl (e.g., guanidinoethyl, guanidinopropyl, guanidinobutyl, guanidinopentyl, and the like) and derivatives thereof; aminobenzyl and derivatives thereof; pyridinium and derivatives thereof; and other like cationic moieties that are known to those of ordinary skill in the art.


Suitable neutral sidechains include, for example, (S) or (R)- α -methylbenzyl and derivatives thereof; benzyl and derivatives thereof; phenethyl and derivatives thereof; naphthylmethyl and derivatives thereof; (S) or (R)- α -methylnaphthyl and derivatives thereof; N-propylpyrrolidinone and derivatives thereof; cyclohexylmethyl and derivatives thereof; furfuryl and derivatives thereof; 3,4,5-trimethoxybenzyl and derivatives thereof; methoxyethyl and derivatives thereof; p-methoxyphenethyl and derivatives thereof; isoamyl ("IsoA") and derivatives thereof; and other like neutral moieties that are known to those of ordinary skill in the art.

Suitable anionic sidechains include, for example, carboxy methyl, carboxy ethyl, and the like, and derivatives thereof; benzoic acid and derivatives thereof; phosphates and derivatives thereof; sulfates and derivatives thereof; and other like anionic moieties that are known to those of ordinary skill in the art.

Optionally, R_1 can be a moiety found on naturally- or non-naturally-occurring amino acids, or R_1 can be a lipid moiety optionally bonded to a linker moiety. As used herein, the term "naturally-occurring amino acid" refers to Ala, Cys, Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr. The term "non-naturally-occurring amino acid" refers to amino acids typically not found in nature, including, for example, D-isomers of naturally-occurring amino acids, 2-aminoadipic acid, 2-aminobutyric acid, norvaline, norleucine, ornithine, and the like.

Typically R_1 is not hydrogen for at least two monomeric units, more typically R_1 is not hydrogen for at least three monomeric units if $n \times m$ is 3 or more. Typically, less than about 75% of the monomer units have an R_1 that is hydrogen. More typically, less than about 50% of the monomer units have an R_1 that is hydrogen. Even more typically,

less than about 25% of the monomer units have an R_1 that is hydrogen. Even more typically R_1 is not hydrogen for any of the monomeric units.

W is typically $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2-$  (i.e., toluic acid), $-\text{CH}_2\text{CH}_2-\text{O}-$, $-\text{CH}_2-\text{CH}=\text{CH}-$, or

5 (II) $-\text{CR}_2\text{R}_3-$,

where R_2 and R_3 for each monomeric unit are monovalent moieties independently selected from a hydrogen atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group; $-\text{SH}$; an optionally substituted, branched or straight chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that
10 optionally contains nitrogen, oxygen, sulfur, phosphorus, and the like, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, phosphorus, and the like; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a
15 backbone structure that optionally contains nitrogen, oxygen, sulfur, phosphorus, and the like, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and a lipid moiety that is optionally bonded to a linker moiety,

wherein when either R_2 and R_3 is an aryl or arylalkyl group having fewer than about 5 carbon atoms in a backbone structure, the backbone structure typically
20 comprises one or more heteroatoms, such as, for example, oxygen and/or nitrogen.

When R_2 and R_3 are aliphatic, they typically contain at least 2 carbon atoms in a backbone structure and more typically contain at least about 3 carbon atoms in a backbone structure. Aryl and arylalkyl R_2 and R_3 groups can be linear or cyclic. Aryl and arylalkyl R_2 and R_3 having less than about 5 carbon atoms in a backbone structure,
25 also typically have one or more heteroatoms in the backbone structure, such as, for example, nitrogen and/or oxygen. Typically aryl and arylalkyl R_2 and R_3 have at least about 5 carbon atoms in a backbone structure.

R_2 and R_3 typically are moieties found on naturally-occurring and non-naturally-occurring amino acids. Usually, at least one of R_2 and R_3 is a hydrogen atom. Most
30 typically, R_2 and R_3 are both hydrogen for all monomeric units, such that compound (I) is a lipid-conjugated, N-substituted polyglycine compound.

The lipid moiety can be positioned at R_a , R_c , R_l for one or more monomers, or at a substitution position in W for one or more monomers. Lipid moieties can be bonded directly to a monomeric unit, or they can be bonded indirectly to a monomeric unit via a linker moiety.

5 The term "linker" used herein refers to a moiety that functions to couple the oligomeric amide and lipid moieties together in a manner such that the molecular distance between the two moieties is greater than would be if the lipid and oligomeric amide moieties were coupled directly to each other. Linker moieties can be relatively small, having from 1 to about 20 atoms in a backbone, or alternatively polymeric. Small
10 linker moieties are optionally substituted and typically have from 1 to about 20 atoms in a backbone (e.g., carbon, nitrogen, oxygen, sulfur, phosphorus, and the like). Typically, small linker moieties have less than about 18 atoms in a backbone, and more typically, less than about 15 atoms in a backbone. Usually, small linker moieties have less than about 10 atoms in a backbone, and optionally have less than about 5 atoms in a
15 backbone.

Linker moieties can be derived from bifunctional molecules such as, for example, 6-aminohexanoic acid, 2-(2-(2-aminoethoxy)ethoxy)ethoxy acetic acid, and the like) that are capable of reacting with both oligomeric and lipid reactants. Linker moieties also can be derived from groups such as, for example, acyl and substituted-acyl groups,
20 sulfonyl and substituted-sulfonyl groups, and other like reactive moieties that are employed during chemical synthesis to facilitate conjugation of the lipid moiety to the oligomeric moiety.

Polymeric linker moieties are optionally substituted (e.g., hydroxy-, carboxy-, phospho-, amino-, and the like), substantially linear polymers having a backbone that
25 contains carbon, and optionally contains nitrogen, oxygen, sulfur, phosphorus, and the like. Polymeric linker moieties have an average molecular weight between about 300 daltons and about 15,000 daltons, typically less than about 10,000 daltons, more typically less than about 5,000 daltons, and even more typically less than about 3000 daltons, and optionally less than about 1000 daltons. Suitable polymeric linker moieties
30 include, for example, polyethylene glycols, polypropylene glycols, polyvinyl alcohols, polyvinylpyrrolidones, and the like.

Lipid moieties are hydrophobic moieties or amphipathic moieties that are either neutral (i.e., having no charge or a net charge of zero) or charged, and either naturally or synthetically derived. Typically, the lipid moiety in lipid-conjugated polyamide compounds of the present invention is amphipathic.

5 Suitable lipid moities include: (1) optionally substituted, aryl or arylalkyl moities having from about 14 to about 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, phosphorus, and the like, where the alkyl component of the arylalkyl moiety optionally has one or more double or triple bonds; (2) optionally substituted, branched or straight chain aliphatic moities having from about 10 to about
10 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, phosphorus, and the like, and optionally has one or more double or triple bonds.

Typically, aryl and arylalkyl lipid moities have at least about 16 carbon atoms and more typically have at least about 20 carbon atoms, and even more typically at least about 30 carbon atoms.

15 Aliphatic lipid moities employed in compounds of the present invention typically have at least about 12 carbon atoms and more typically have at least about 14 carbon atoms. Usually, the aliphatic lipid moities have at least about 18 carbon atoms, more usually at least about 24 carbon atoms, and even more usually at least about 30 carbon atoms.

20 The number of lipid moities in lipid-conjugated polyamide compounds of the present invention can vary depending on the degree of hydrophobicity desired, and will also vary with oligomer length (i.e., $n \times m$) and size of lipid moiety. For example, when the lipid moiety has about 30 carbon atoms or less, lipid-conjugated polyamide compounds of the present invention typically have conjugated to it, a number of lipid
25 moities that is less than the number computed as 90% of the total number of monomeric groups (i.e., $n \times m$) (i.e., if n is 3 and m is 3, then the number of lipid moities conjugated to the lipid-conjugated polyamide compound is typically less than about 8). More typically, when the lipid moiety has about 30 carbon atoms or less, lipid-conjugated polyamide compounds of the present invention have conjugated to it, a number of lipid
30 moities that is less than about 80% of the total number of monomeric groups. more

typically less than about 75% of the total number of monomeric groups, and even more typically less than about 60% of the total number of monomeric groups.

When the lipid moiety has more than about 30 carbon atoms, typically, lipid-conjugated polyamide compounds of the present invention have conjugated to it a
5 number of lipid moieties that is less than the number computed as 50% of the total number of monomeric groups.

Suitable lipid moieties include those having one or more hydrophobic tails that are optionally substituted aliphatic, straight chain moieties, each hydrophobic tail independently having from about 8 to about 30 carbon atoms in a backbone that in
10 addition, optionally contains nitrogen, oxygen, sulfur, phosphorus, and the like. Typically, hydrophobic tails have at least about 10 carbon atoms in a backbone and more typically have at least about 12 carbon atoms in a backbone. Hydrophobic tails employed in lipid-conjugated polyamide compounds of the present invention typically do not have more than about 26 carbon atoms in a backbone, and more typically do not
15 have more than about 24 carbon atoms in a backbone.


Natural lipid moieties employed in the practice of the present invention can be derived from, for example, phospholipids, including, for example, phosphoglycerides (including both acyl phosphoglycerides (such as, for example, phosphatidic acid, phosphatidyl ethanolamine, phosphatidyl serine, phosphatidyl inositol, phosphatidyl
20 inositol phosphate, phosphatidyl inositol biphosphate, phosphatidyl glycerol, diphosphatidylglycerol, and the like) and ether phosphoglycerides); glycosylglycerides (such as, for example, monogalactosyl diacylglycerol, digalactosyldiacylglycerol, sulphoquinovosyldiacylglycerol, dimannosyldiacylglycerol, galactofuranosyldiacylglycerol, galactosylglucosyldiacylglycerol,
25 galactosylglucosyldiacylglycerol, glucosylgalactosylglucosyldiacylglycerol, and the like); sphingolipids (such as, for example, sphingosines, glycosyl ceramides, gangliosides, and the like); and saturated and unsaturated sterols (such as, for example, cholesterol, ergosterol, stigmasterol, sitosterol, and the like); and other like natural lipids.

30 Suitable synthetic lipid moieties can be derived from, for example, dipalmitoyl phosphatidylethanolamine (DMPE) (Genzyme Corp., Cambridge), DMR1E-C™

(GibcoBRL, Gaithersburg, MD), 2,3-dioleoyloxy-N-[2(spermine-carboxamido)ethyl]-N,N-dimethyl-1-propanaminiumtrifluoroacetate (DOSPA) (Lipofectamine™, GibcoBRL, Gaithersburg, MD), 3β-[N-(N',N'-dimethylaminoethyl)carbonyl]cholesterol, Tfx-50 (Promega Corp., Madison, WI), N,N1,N2,N3-tetramethyl-N,N1,N2,N3-tetrapalmitylsperimine (TM-TPS) (Cellfectin, GibcoBRL, Gaithersburg, MD), dipalmitoyl phosphatidylethanolaminospermine, and the like.

Suitable lipid moieties also include those derived from fatty acids and fatty alcohols having from about 8 to about 24 carbon atoms in a backbone. Typically, the fatty acids and fatty alcohols have at least about 10 carbon atoms in a backbone, and more typically have at least about 12 carbon atoms in a backbone. Usually, the fatty acids and alcohols from which lipid moieties are derived have less than about 20 carbon atoms in a backbone.

Typically, R_a is a lipid moiety or a lipid moiety conjugated to a linker moiety. A particularly useful lipid moiety-containing R_a radical is the phosphatidyl alkylamino-substituted acyl radical having the formula,




where p is an integer selected from 2 or 3, and each R_4 is independently selected from an alkyl or alkenyl moiety having from about 6 and to about 25 carbon atoms in a backbone. Typically R_4 has up to about 22 carbon atoms in a backbone, more typically, up to about 20 carbon atoms, even more typically up to about 18 atoms. Typically, R_4 has at least about 8 carbon atoms in a backbone, more typically at least about 10 carbon atoms, and even more typically at least about 12 carbon atoms in a backbone. Exemplary R_4 moieties include dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, and the like. Preferably p is 2.

Lipid-conjugated polyamide compounds of the present invention can be optionally further conjugated or complexed with agents that impart, for example, targeting capabilities, structural features, biological activity, or that introduce

degradations sites, and the like. Suitable agents include, for example, mono-, di-, and polysaccharides, polyethylene glycols, amino acids, peptides, polypeptides, proteins (including, for example, lipoproteins, glycoproteins, antibodies, and the like), crosslinking agents, marker agents (such as, for example, fluorescein, biotin, ^{32}P , and the like), and the like.

Those of ordinary skill in the art will appreciate that R_1 , R_c , R_a , W , and the particular lipid moiety employed can be readily varied to optimize the physicochemical properties of the lipid-conjugated polyamide compound for delivery of a particular type of biologically active compound. For example, oligomeric moieties of the present invention suitable for use in the delivery of polynucleic acids to cells have a net positive charge and are capable of condensing polynucleic acids so that they are more compact in size, thus facilitating their delivery to cells.

Compounds of formula (I) that are suitable for use in the delivery of polynucleic acids to cells, include lipid-conjugated polyamide compounds having repeating n -mer units (i.e., where n is greater than 1). For example, when n is 3, the lipid-conjugated polyamide compound of formula (I) has repeating trimer units, i.e.,



where R_a , R_c , m , each W and each R_1 are defined as in formula (I). Compounds having formula (IV) that are suitable for use in the delivery of polynucleic acids to cells include, for example, those where R_1^1 is a cationic side chain, R_1^2 and R_1^3 are both neutral side chains, each W is CH_2 , R_c is NH_2 , and R_a is defined by formula (III), such as those compounds shown in Table 2, in Example 1, herein.

Lipid-conjugated polyamide compounds of the present invention typically form concentration-dependent, ordered two- or three-dimensional structures in solution. Such structures include two dimensional arrays, such as, for example, a single charged layer or a lipid bilayer surface, and three-dimensional structures, such as, for example, micelles, vesicles, and liposomes. Typically, ordered structures formed from lipid-conjugated polyamide compounds of the present invention by themselves, typically are sufficiently small such that they do not scatter light. Micelles, vesicles, and liposomes prepared from

lipid-conjugated compounds complexed with polynucleotides typically have average particle sizes that are less than about 1 μm , more typically less than about 500 nm, and even more typically less than about 200 nm.


In addition to the delivery of biologically active agents to cells, lipid-conjugated polyamide compounds of the present invention can also be used in applications, such as, for example, screening peptide-like compounds for biological activity, incorporation into biosensors such that the oligomeric moiety has the capacity to bind to a target ligand, and the like. For drug screening applications, for example, libraries of lipid-conjugated polyamide compounds having a variety of R_1 groups can be synthesized and subsequently screened for biological activity in accordance with the methods for synthesizing and screening modified peptide libraries described in PCT publication WO 91/19735 (published December 26, 1991), incorporated herein by reference.

SYNTHESIS OF LIPID-CONJUGATED POLYAMIDE COMPOUNDS

Lipid-conjugated polyamide compounds of the present invention can be synthesized by both solid-phase and solution-phase methods. The present invention also provides a method of synthesizing lipid-conjugated polyamide compounds, said method comprising:

a) contacting

- (1) a lipid reactant, with
- (2) an oligomer reactant, wherein said oligomer reactant has the general formula:



wherein n is an integer selected from 1 to about 48, and m is an integer from about 2 to about 48,

wherein each T_a and T_c is independently selected from a terminal group and a reactive moiety that is capable of further reaction with said lipid reactant,

wherein R_1 for each monomeric unit, $\text{---} \begin{array}{c} R_1 \\ | \\ \text{N} - \text{W} - \text{C} \\ || \\ \text{O} \end{array} \text{---}$ in said oligomer reactant is selected from the group consisting of a hydrogen atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group, -SH; an optionally substituted, branched or straight chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and a reactive moiety that is capable of further reaction with said lipid reactant,

wherein when R_1 , R_a , or R_c is an aryl or arylalkyl group having fewer than about 5 carbon atoms in a backbone structure, the backbone structure typically comprises one or more heteroatoms, such as, for example, oxygen and/or nitrogen,

wherein R_1 is not a hydrogen atom for at least one monomeric unit,

wherein W for each monomeric unit is selected from an optionally substituted, branched or straight chain divalent moiety having from 1 to about 50 atoms in a backbone that contains carbon, and optionally contains nitrogen, oxygen, sulfur, and phosphorus, and optionally one or more double or triple bonds, wherein said optional substitution of W may be a reactive moiety that is capable of further reaction with said lipid reactant,

wherein at least one of T_a , T_c , W for a single monomeric unit, or R_1 for a single monomeric unit comprises a reactive moiety that is capable of further reaction with said lipid reactant; then

b) reacting said lipid reactant with said oligomer reactant to conjugate the lipid reactant to the oligomer reactant.

The term "lipid reactant" used herein refers to a reactive species having a lipid moiety that is capable of participating in a chemical reaction, such as, for example, nucleophilic displacement, condensation, and the like. Lipid reactants having functional groups, such as, for example, $-NH_2$, $-COOH$, $-SH$, $-OH$, $-SO_2Cl$, and $-CHO$ are particularly useful for synthesizing lipid-conjugated compounds of the present invention. Lipid reactants suitable for use in the practice of the present invention include lipid reactants having any one of the lipid moieties described herein which can react with, or which can be modified to react with, the oligomeric reactant or a linker. Typically, lipid reactants are primary, secondary, or tertiary amines. Preferred lipid reactants are phosphatidylethanolamines.

As used herein, the term "oligomer reactant" refers to an oligomeric amide that is capable of participating in a chemical reaction, such as, for example, nucleophilic displacement, condensation, and the like. Oligomer reactants typically are acylated with a leaving group that is susceptible to nucleophilic displacement by a nucleophile, such as an amine. Oligomer reactants suitable for use in the practice of the present invention include all of the oligomeric amide substituents described for formula (I)

(i.e., $\text{---}[(\overset{R_1}{\underset{|}{N}}\text{---}W\text{---}\overset{O}{\underset{||}{C}})]_n\text{---}$) herein.

As used herein, the term "reactive moiety" refers to a moiety that is capable of participating in a reaction with the lipid reactant. Typical reactive moieties include, for example, $-NH_2$, $-OH$, $-H$, $-SH$, $-COOH$, acyl (e.g., acetyl), benzoyl, sulfonyl (e.g., dansyl), amide, hydrazine (typically a T_c group), and derivatives thereof (including alkyl-substituted derivatives), and the like. Typically, the reactive moiety is an acyl moiety substituted with a leaving group that is susceptible to nucleophilic displacement by a nucleophile, such as an amine.

Exemplary terminal groups include R_a and R_c moieties that are not reactive moieties, moieties that are biologically active agents, targeting agents (e.g., a cell receptor ligand, antibody, etc.), marker agents, amino acid residues that function, for

example, as a degradation site for endogenous proteolytic enzymes, and the like. These terminal groups typically are not further reactive with the lipid reactant.

The oligomer reactant and lipid reactant can be optionally bonded to each other via a linker moiety (which optionally can be derived from a reactive moiety).

5 Alternatively, the linker moiety, which is derived from a molecule that is capable of reacting with both oligomeric and lipid reactants, can be optionally conjugated to either the lipid or oligomer reactant prior to reaction between lipid and oligomer reactants. Thus, the lipid reactant can be conjugated to the oligomer reactant either directly, or indirectly via the linker moiety.

10 The term "reacting" used herein refers to one or more chemical reactions that result in formation of a chemical bond between the lipid reactant and the oligomer reactant, either directly, or indirectly via the linker moiety. Suitable reactions include, for example, condensation (e.g., acylation, and the like) and nucleophilic displacement.

Oligomer reactants having the general formula (IV) can be prepared, for example,
15 via a series of nucleophilic displacement reactions according to the solid-phase method described by Zuckermann et al., PCT WO94/06451 (published March 31, 1994), incorporated herein by reference. The method can be performed utilizing automated peptide synthesis instrumentation to permit rapid synthesis of oligomer reactants of interest. These instruments are commercially available from, for example, Applied
20 Biosystems.

Specifically, monomer assembly into oligomer reactants is achieved by the sequential addition of "submonomer" units to the growing chain. In one method of monomer assembly, each cycle of monomer addition consists of two steps:

(1) acylation of a secondary amine bound to the solid support with an
25 acylating agent that has a leaving group (i.e., a group susceptible to nucleophilic displacement by a nucleophile, such as an amine) and a carbonyl group (e.g., a carboxyl group) (i.e., the "acylation step"); followed by

(2) nucleophilic displacement of the leaving group with a sufficient amount
of a submonomer that has a primary, secondary, or tertiary amino group to introduce a
30 side-chain (i.e., the "nucleophilic displacement step").

A schematic of the reaction scheme is shown in Figure 1. Exemplary acylating agents include haloacetic acid, halomethyl benzoic acid, and the like. The efficiency of displacement of the leaving group is modulated by the type of acylating agent employed. For example, when a haloacetic acid is employed, it has been observed that iodine is more efficient at displacing the leaving group compared to chlorine. Suitable submonomers include alkylamines, alkenylamines, aromatic amines, alkoxyamines, semicarbazides, acyl hydrazides, and derivatives thereof, and the like.

Oligomer synthesis using the submonomer approach occurs in the carboxy to amino direction. The oligomer is elaborated until the desired length, then is terminated, for example, with a bromoacetamide group. One advantage of using solid phase submonomer assembly to construct oligomer reactants of the present invention is that the need for N- α -protected monomers is eliminated, as only reactive side-chain functionalities need to be protected.

Typically, the oligomeric reactant is synthesized as a series of repeating di-, tri-, or tetra-mer units. An exemplary trimer-based cationic oligomer has the following monomer sequence in the amino terminal (T_a) to carboxy terminus (T_c) direction:

- (1) positively charge monomer
- (2) neutral monomer, and
- (3) neutral monomer.

The terms "neutral monomer" and "positively charged monomer" as used herein refer to the net charge of the monomeric unit.

Further reaction of the oligomer reactant with the lipid reactant can occur by further acylation and/or nucleophilic displacement. For example, an oligomer reactant that is haloacylated (e.g., where T_a is a bromoacetyl group) can be reacted with an lipid reactant that is a primary, secondary, or tertiary amine. Conjugation thus occurs by nucleophilic displacement of the bromine, to form a lipid-conjugated polyamide compound.

LIPID-CONJUGATED POLYAMIDE COMPOSITIONS AND USES THEREOF

In yet another embodiment, the present invention provides compositions comprising lipid-conjugated polyamide compound(s) of the present invention and an

effective amount of a biologically active agent. As used herein, the terms "effective amount" and "effective dose" refer to an amount of biologically active agent or biologically active agent-containing lipid-conjugated polyamide composition of the present invention that is sufficient to detectably induce, or participate in, a biological response, such as, for example, signal transduction, transcription, translation, lymphocyte activation, including, for example, antibody production, and the like, in for example, a cell, a mammal, or a bird. The term "biologically active agent" used herein refers to an agent that upon administration in an effective amount to, for example, a cell, a mammal, or a bird, induces or participates in, a biological response.

When the biologically active agent is a polynucleotide, the relative quantities of lipid-conjugated polyamide compound to polynucleic acid are typically selected such that the +/- charge ratio of lipid-conjugated polyamide compound to polynucleotide in the composition is at least about 2 and less than about 10. More typically, the +/- charge ratio is less than about 8, and even more typically is less than about 4. The charge ratio is computed according to the following:

$$\text{Charge Ratio} = (n_L \times M_L) / (3.03 \times M_{\text{DNA}}),$$

where n_L is the number of moles of lipid-conjugated polyamide compound, M_L = net number of charges/mole lipid-conjugated polyamide, and where M_{DNA} = micrograms of DNA.

Compositions of the present invention can be in liquid or solid form, and can optionally include pharmaceutically acceptable excipients. Such excipients can be used as fillers, processing aids, delivery enhancers and modifiers, and the like. Suitable excipients include, for example, calcium phosphate, magnesium stearate, talc, monosaccharides, disaccharides, polysaccharides, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, dextrose, polyvinyl alcohol, polyvinylpyrrolidone, low melting waxes, ion exchange resins, and the like, as well as combinations of any two or more thereof. A thorough discussion of pharmaceutically acceptable excipients is available in "Remington's Pharmaceutical Sciences" (Mack Pub. Co., NJ 1991), which is incorporated herein by reference.

Additional agents can be included in the compositions, such as, for example, marker agents, nutrients, and the like. For example, when the biologically active agent is

a polynucleotide, agents that promote endocytosis of the desired nucleic acids or aid in binding of the nucleic acids to the cell surface, or both, can be incorporated into compositions of the present invention.

Liquid compositions of the present invention can be in the form of a solution,
5 suspension, or emulsion with a liquid carrier. Suitable liquid carriers include, for example, water, saline, pharmaceutically acceptable organic solvent(s), pharmaceutically acceptable oils or fats, mixtures thereof, and the like. The liquid carrier may contain other suitable pharmaceutically acceptable additives, such as solubilizers, emulsifiers, nutrients, buffers, preservatives, suspending agents, thickening agents,
10 viscosity regulators, stabilizers, and the like. Suitable organic solvents include, for example, monohydric alcohols, such as ethanol, and polyhydric alcohols, such as glycols. Suitable oils include, for example, soybean oil, coconut oil, olive oil, safflower oil, cottonseed oil, and the like. For parenteral administration, the carrier can also be an oily ester such as ethyl oleate, isopropyl myristate, and the like.

15 In yet another embodiment, the present invention provides a method for inducing the uptake of a biologically active agent by a cell, said method comprising:

providing a composition comprising an effective amount of a biologically active agent and a lipid-conjugated polyamide compound; then

20 contacting a biological sample with an effective dose of said composition, wherein said biological sample comprises a cell.

As used herein, the term "biological sample" refers to a sample comprising one or more cells or tissue. Cells suitable for use in the practice of the present invention include, for example, mammalian cell lines available from the American Type Culture Collection (ATCC), Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster
25 kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), human embryonic kidney cells, baby hamster kidney cells, mouse sertoli cells, canine kidney cells, buffalo rat liver cells, human lung cells, human liver cells, mouse mammary tumor cells, other mammalian (including human) cells (e.g., stem cells, particularly hemapoitic cells, lymphocytes, macrophages, dendritic cells, tumor cells and
30 the like), and the like.

Suitable tissue for use as samples in the present invention include, for example, tissue derived from mammals, such as, muscle, skin, brain, lung, liver, spleen, blood, bone marrow, thymus, heart, lymph, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, connective,
5 and the like.

Modes of administration to a sample include, for example, *ex vivo* administration to samples derived from a subject and *in vitro* administration to a sample. Methods for carrying out these modes of administration are well known to those of ordinary skill in the art. For example, when the biological agent is a polynucleotide, *ex vivo* delivery and
10 reimplantation of transformed cells into a subject can be achieved as described in e.g., International Publication No. WO 93/14778 (published August 5, 1993), which is incorporated herein by reference.

In yet another embodiment, the present invention provides a method for inducing the uptake of a biologically active agent by a cell *in vivo*, said method comprising:
15 providing a composition comprising an effective amount of a biologically active agent and a lipid-conjugated polyamide compound; then administering an effective dose of said composition to a subject.

The present invention further provides a method of expressing a gene in a mammal, said method comprising:
20 administering a polynucleic acid complexed with a lipid-conjugated polyamide of the present invention to a mammal, wherein said polynucleic acid is capable of functionally expressing said gene in said mammal, and wherein said complex is effective at transfecting said gene into a cell in
25 said mammal.

As used herein, the term "subject" refers to birds and mammals, including for example, rodents and humans. Direct administration to a subject can typically be accomplished by injection, either subcutaneously, intraperitoneally, intravenously or intramuscularly or delivered to the interstitial space of a tissue. The compositions can
30 also be administered into a tumor or lesion. Other modes of administration include oral

and pulmonary administration, suppositories, and transdermal applications, and the like, as well as administration via needles, and gene guns or hyposprays.

Lipid-conjugated compounds of the present invention, when combined with a biologically active agent, are effective at inducing the uptake the biologically active agent by cells. As used herein, the term "induce" and its various grammatical equivalents, refers to effecting the uptake of a biologically active agent by a cell. Methods used for detecting the uptake of biologically active agents by cells will vary depending on the type of biologically active agent employed, however, those of ordinary skill in the art will appreciate that cell uptake can be detected by a variety of known assays and histological techniques, as well as by a variety of diagnostic methods, including, for example, clinical diagnostic methods (e.g., alleviation of symptoms, etc.).

Typically, compounds of the present invention are effective at enhancing the uptake the biologically active agent by cells. When uptake of a biologically active agent by a cell is enhanced, typically the uptake of biologically active agent by the cell is at least about 5% greater than the uptake of the biological agent in neat form (e.g., substantially free of lipid-conjugated polyamide compound). More typically, when uptake is enhanced, the uptake of the biologically active agent by the cell is at least about 10% greater than the uptake of the biologically active agent in neat form, even more typically at least about 15% greater, and even more typically at least about 20% greater.

An effective amount of biologically active agent and likewise, an effective dose of lipid-conjugated polyamide/biological agent containing composition will, of course, vary depending upon known factors such as the pharmacodynamic characteristics of the particular biologically active agent, its mode and route of administration; the age, health, and weight of the recipient; nature and extent of symptoms; kind(s) of concurrent treatment, frequency of treatment, the effect desired, and the like. However, the precise amount for a particular patient and biologically active agent can be readily determined by routine experimentation by a clinician of ordinary skill in the art. Those having ordinary skill in the art will also appreciate that the precise amount of lipid-conjugated compound employed in compositions of the present invention can be readily determined by routine screening studies to determine the optimal ratio of lipid-conjugated polyamide compound to biologically active agent for inducing the desired response. Compositions

of lipid-conjugated polyamide compounds complexed with biologically active agents can be administered as a single dose or in multiple doses. Multiple doses can be administered either continuously, in intervals, or a combination of both. For purposes of the present invention, an effective *in vivo* amount will be from about 0.01 mg/kg/day to about 50 mg/kg/day or about 0.05 mg/kg/day to about 10 mg/kg/day of biologically active agent.

Compositions of the present invention are particularly effective at inducing the uptake of polynucleotides into cells. Polynucleotide uptake by cells can be detected by using protein expression assays or polynucleotide hybridization techniques.

Compositions can be screened and optimized with respect to transfection efficiency by incorporating a reporter gene into the DNA and assaying for the reporter gene product using standard immunoassay methods or biological or enzymatic activity assays (such as, for example, a luciferase assay).

When combined with polynucleotides, lipid-conjugated polyamide compounds are particularly effective at inhibiting nuclease-induced polynucleotide degradation caused by nucleases present in serum, and other biological fluids. As a result, smaller quantities of polynucleotides can be more efficiently administered using compositions of the present invention.

Thus, the present invention also provides a method for substantially inhibiting nuclease-induced polynucleotide degradation, said method comprising contacting a polynucleotide with a degradation-inhibiting quantity of lipid-conjugated polyamide compound, and introducing the polynucleotide and the lipid-conjugated polyamide compound into a nuclease-containing environment. Inhibition of nuclease-induced polynucleotide degradation can be detected by incubating a sample of lipid-conjugated polyamide protected polynucleotide and a control sample of unprotected polynucleotide in, for example, serum, for various time periods, then analyzing the mixtures by gel electrophoresis to determine when degradation occurs (for example, when about 50% of the polynucleotide has degraded) for the protected polynucleotide, as compared to unprotected polynucleotide. The degree of degradation can be monitored using known quantitation methods, such as, for example, radiolabelling and the like.

Typically, a degradation-inhibiting quantity of lipid-conjugated polyamide compound is the amount of lipid-conjugated polyamide compound sufficient to substantially inhibit nuclease-induced polynucleotide degradation of a polynucleotide for at least about 5 minutes, as compared to unprotected (i.e., neat) polynucleotide. More typically, the quantity of lipid-conjugated polyamide compound is sufficient to inhibit nuclease-induced polynucleotide degradation for at least 10 minutes, more typically for at least about 30 minutes, more typically for at least 60 minutes, more typically for at least 90 minutes, and even more typically for at least 120 minutes, as compared to unprotected polynucleotide. Lipid-conjugated polyamide compounds of the present invention are effective at inhibiting nuclease-induced polynucleotide degradation to the extent that lipid-conjugated polyamide-protected polynucleic acid typically exhibit less than about 80% of the degradation exhibited by unprotected (i.e., neat) polynucleotide during the same time period, more typically less than about 50%, even more typically less than about 30%, and more typically, less than about 20%.

Compositions of lipid-conjugated polyamide compounds and polynucleic acid that are particularly effective at enhancing polynucleotide uptake by cells are those where the lipid-conjugated polyamide compound has formula (I), where W is CH₂, R_a contains a lipid moiety (preferably a phosphatidylethanolamine), n is greater than 1, and each n-mer contains both cationic and neutral R₁ sidechain groups. This lipid-conjugated polyamide compound is also effective at protecting polynucleotides from nuclease-induced degradation.

DILUTION-CONCENTRATION METHOD FOR MAKING STABLE POLYNUCLEIC ACID/DELIVERY VEHICLE COMPLEX PREPARATIONS

In still a further embodiment, the present invention provides a method of making a stable preparation of a polynucleic acid complexed with a delivery vehicle. said method comprising:

- a) combining a polynucleic acid with a first liquid carrier to form a dilute polynucleic acid solution that is substantially precipitant-free;
- b) combining a delivery vehicle-forming compound with a second liquid carrier to form a delivery vehicle solution that is substantially precipitant-free;

c) combining said dilute polynucleic acid solution with said delivery vehicle solution to form a dilute preparation of delivery vehicle/polynucleic acid complex; then

d) reducing the volume of said dilute preparation to form a stable preparation of delivery vehicle/polynucleic acid complex,

wherein the concentration of polynucleic acid in said stable preparation is higher than the concentration of polynucleic acid in said dilute polynucleic acid solution, and

wherein said stable preparation is substantially precipitant-free.

It has been discovered that when delivery vehicle/polynucleic acid complexes are formulated in liquid carrier in accordance with the "dilution-concentration" method of formulation, the resulting complexes are stable for relatively long periods of time with respect to particle size and transfection efficiency. In addition, the stable preparations are characteristically precipitant-free. Thus, unlike conventional delivery vehicle/polynucleic acid preparations in which equal volumes of dilute solutions of polynucleic acid and delivery vehicle are mixed together just prior to transfection, stable preparations made according to the invention dilution-concentration method of formulation can be prepared and stored up to days prior to transfection without substantial change in transfection efficiency. As used herein, the term "transfection efficiency" refers to the quantity or activity of protein expressed, or polynucleic acid synthesized, as a result of administration of a delivery vehicle/polynucleotide complex to a cell.

Typically, the particle size distribution of delivery vehicle/polynucleotide complexes in the stable preparations changes by less than about 40% one day after formulation by the invention method, as compared to the particle size distribution of the complexes immediately after formulation. More typically, the particle size distribution of delivery vehicle/polynucleotide complexes in the stable preparations changes by less than about 30%, more typically by less than about 20%, and even more typically by less than about 15%, one day after formulation according to the concentration method of the present invention, as compared to the particle size distribution of the complexes immediately after formulation.

More typically, the particle size distribution of the delivery vehicle/polynucleotide complexes in the stable preparations changes by less than about 40%, even more typically less than about 30%, usually less than 20%, and more usually less than about 15%, five to eight days after formulation by the invention method, as
5 compared to the particle size distribution of the complexes immediately after formulation.

As used herein, the term "dilute polynucleic acid solution" refers to a solution having a concentration of less than about 300 μg polynucleic acid/ml. Typically, dilute polynucleic acid solutions have polynucleic acid concentrations of less than about 250
10 $\mu\text{g}/\text{ml}$, more typically less than about 150 $\mu\text{g}/\text{ml}$, and even more typically less than about 100 $\mu\text{g}/\text{ml}$. Preferably, dilute polynucleic acid solutions have polynucleic acid concentrations of less than about 50 $\mu\text{g}/\text{ml}$.

The concentration of polynucleic acid in stable preparations of delivery vehicle/polynucleic complex is greater than the polynucleic acid concentration of the
15 dilute polynucleic acid solution, and is typically at least about 150 μg (polynucleotide)/ml (preparation). More typically, the concentration of polynucleotide in stable preparation is at least about 250 $\mu\text{g}/\text{ml}$, more typically at least about 500 $\mu\text{g}/\text{ml}$, and even more typically at least about 1 mg/ml, and even more typically at least about 2 mg/ml

20 The concentration of delivery vehicle-forming compound employed in the delivery vehicle solution will vary depending on the ratio of DNA to delivery vehicle desired and the solubility properties of the delivery-vehicle-forming compound employed. Those having ordinary skill in the art will recognize that the concentration and volume of delivery vehicle-forming compound in second liquid carrier can be
25 adjusted to achieve the desired target ratio of delivery vehicle to DNA.

When the delivery vehicle-forming compound is a lipid-conjugated polyamide of the present invention, the concentration of the delivery vehicle solution is typically not more than about 5 mg/ml, more typically not more than about 2.5 mg/ml, and typically at least about 1.25 mg/ml. The volumes of delivery vehicle solution and dilute polynucleic
30 acid solution combined are preferably selected such that the ratio of lipid-conjugated polyamide to polynucleic acid is about 5 mg lipid-conjugated polyamide to about 1 mg

polynucleic acid. Thus, for example, if a 5 mg/ml lipid-conjugated polyamide solution is used as the delivery vehicle solution, combination with an equal volume of a 1 mg/ml polynucleic acid solution will yield a ratio of lipid-conjugated polyamide to polynucleic acid of 5 mg lipid-conjugated polyamide to 5 mg polynucleic acid.

5 The first liquid carrier may be the same or different from the second liquid carrier. Suitable first and second liquid carriers include the liquid carriers described herein. The term "delivery vehicle" used herein refers to an ordered structure that is capable of complexing with or enveloping a polynucleotide. Suitable delivery vehicles include liposomes, micelles, vesicles, and the like. As used herein, the term "delivery
10 vehicle-forming compound" refers to a compound that is capable of forming a delivery vehicle. Exemplary delivery vehicle-forming compounds include amphipathic lipids, non-lipid polycationic compounds, lipid-conjugated polyamide compounds, and the like.

 After the dilute polynucleotide solution and delivery vehicle preparation are combined, the volume of the resulting mixture is reduced. Suitable methods for reducing
15 the volume of the resulting mixture include, for example, centrifugal filtration, vacuum filtration, as well as other methods for volume reduction that are well known to those having ordinary skill in the art. Typically, the volume of the resulting mixture is reduced such that the concentration of polynucleotide in the preparation is greater than about 150 µg/ml. The final concentration of the concentrated preparation can be lowered after the
20 volume reduction step by adding sufficient liquid carrier to the preparation to achieve the desired concentration.

 As used herein, the term "precipitant-free" refers to a preparation or solution that is free of solid material, i.e., precipitant material, as visually detected by the naked eye.

 It has been discovered that transfection of cells using stable delivery
25 vehicle/polynucleotide complex preparations is typically more efficient than with complexes formed by the conventional method of mixing equal volumes of dilute solutions of polynucleotide and delivery vehicle together, immediately prior to transfection, as illustrated in Figure 14 .

 The invention will now be described in greater detail with reference to the
30 following non-limiting examples.

Example 1Preparation of Lipid-Conjugated Polyamide CompoundsA. Synthesis of Lipid-Conjugated Polyamide Compounds5 (1) Synthesis of Oligomer Reactant

A fritted reaction vessel was charged with 100 mg of Fmoc-Rink amide resin with a substitution level of about 0.50 mmol/g resin. Two milliliters of dimethylformamide (DMF) was added to the resin. The mixture was agitated for 1-2 minutes to swell the resin, then the DMF was drained. The Fmoc groups were removed
10 by adding 2.0 ml of 20% piperidine in DMF to the resin, then agitating for 1 minute. The solution was then drained from the resin. Another 2 ml of 20% piperidine in DMF was added to the resin, followed by agitation for 15 minutes, after which, the DMF was then drained from the resin.

The resin was washed by adding 2 ml of DMF to the resin, followed by agitation
15 to form a uniform slurry. The resin slurry was agitated by bubbling argon up through the bottom of the fritted vessel. The DMF was removed from the resin by vacuum filtration through the fritted bottom of the reaction vessel until the resin appeared dry (typically about 5 seconds). The washing step was repeated 6 times with DMF.

The deblocked amine was then acylated by adding 850 μ l of 1.2 M bromoacetic
20 acid in DMF to the resin, followed by 200 μ l of N,N'-diisopropylcarbodiimide (DIC). This mixture was agitated for 45 minutes at 35°C, then the solution was drained from the resin. The resin was washed with 6 x 2 ml aliquots of DMF. Amine displacement was effected by treating the resin with 850 μ l of a 1 M solution of an amine in dimethylsulfoxide (DMSO). This mixture was agitated at 35 °C for 20 minutes, then the
25 amine solution was drained from the resin. After the amine displacement step, the resin was washed with 6 x 2 ml aliquots of DMF. This completed one cycle of acylation and displacement.

The second cycle was initiated by acylation using bromoacetic acid and DIC, followed by displacement with a second amine as described above. This
30 acylation/displacement cycle was repeated until the desired oligomer was reached.

The N-terminus of the resulting oligomer was acylated at 35°C for 45 minutes with 850 µl of bromoacetic acid in DMF (1.2 M) and 200 µl DIC.

(2) Displacement of Bromide from Bromoacetylated Oligomer Reactant with Dimyristoyl Phosphatidylethanolamine

5 A quantity of 254 mg of dimyristoyl phosphatidylethanolamine (DMPE) (Genzyme, Cambridge, MA) was suspended in 2 ml of 15% methanol in chlorobenzene, then treated with 30 µl of 12.8 N aqueous KOH. The resulting solution of DMPE was added to the bromoacetylated oligomer on resin from (1), above, and the reaction was agitated by bubbling argon for 1 second every 30 seconds over a period of 14 hours at
10 35°C. Next, the resin was thoroughly washed with 15% methanol in chlorobenzene to remove unreacted DMPE.

The resulting lipid-conjugated polyamide compound was cleaved from the resin and deprotected using 95% TFA in water, then lyophilized to yield a crude product.

This method was repeated using dipalmitoyl phosphatidylethanolamine (DPPE)
15 and dioleoyl phosphatidylethanolamine (DOPE). Lipid-conjugated polyamide compounds prepared in accordance with this method are shown in Table 1 and Table 2 below.

Table 1. Lipid-Conjugated N-Substituted Polyglycine Compounds[†]

Compound No.	n	m	R ₁	R ₄
1	1	2	2-aminoethyl	tridecyl
2	1	4	2-aminoethyl	tridecyl
3	1	10	2-aminoethyl	tridecyl
4	1	2	2-aminoethyl	pentadecyl
5	1	4	2-aminoethyl	pentadecyl
6	1	10	2-aminoethyl	tridecyl
7	1	2	3-aminopropyl	tridecyl
8	1	4	3-aminopropyl	tridecyl
9	1	10	3-aminopropyl	tridecyl
10	1	2	3-aminopropyl	pentadecyl
11	1	4	3-aminopropyl	pentadecyl
12	1	10	3-aminopropyl	pentadecyl

[†] Compounds of formula (I) where W is CH₂, R_c is NH₂, and R_s is defined by formula (III), where p is 2.

Table 2. Examples of Lipid-Conjugated N-substituted Polyglycine Compounds Having Repeating Trimer Units[†]

Compound No.	m	R ¹ ₁	R ² ₁ and R ³ ₁	R ₄
13	3	2-aminoethyl	2-phenylethyl	tridecyl
14	3	2-aminoethyl	(S)-1-methylbenzyl	tridecyl
15	2	2-aminoethyl	2-(4'-methoxyphenyl)ethyl	tridecyl
16	3	2-aminoethyl	2-(4'-methoxyphenyl)ethyl	tridecyl
17	4	2-aminoethyl	2-(4'-methoxyphenyl)ethyl	tridecyl
18	8	2-aminoethyl	2-(4'-methoxyphenyl)ethyl	tridecyl
19	12	2-aminoethyl	2-(4'-methoxyphenyl)ethyl	tridecyl
20	3	2-aminoethyl	2-(4'-methoxyphenyl)ethyl	<i>cis</i> -8-heptadecyl
21	3	2-aminoethyl	pentyl	tridecyl
22	3	2-aminoethyl	pentyl	<i>cis</i> -8-heptadecyl
23	3	2-aminoethyl	3-methylbutyl	tridecyl
24	3	2-aminoethyl	3-methylbutyl	<i>cis</i> -8-heptadecyl

5 [†]Compounds of formula (IV) where W is CH₂, R_c is NH₂, and R_s is defined by formula (III), where p is 2.

B. Characterization of Lipid-Conjugated Polyamide Compounds

10 The lyophilized crude product from part A was dissolved in 5 ml of 50% (v/v) acetonitrile in water, and resulting solution was applied to a Delta-Pak™ C4 column (15 μm 300 Å) equilibrated with 80% (v/v) solvent B (0.1% (v/v) TFA in acetonitrile) in solvent A (0.1% (v/v) TFA in water) on a Water Prep LC3000 system equipped with a UV detector (Waters Corp., Milford, MA). Peaks were eluted with a linear gradient of 80% (v/v) solvent B in solvent A to 100% solvent B over 20 minutes, followed by 100% solvent B for 15 minutes at a flow rate of 50 ml/min. Peaks were monitored at 220 nm. the fractions containing the desired lipid-conjugated polyamide compounds were combined and concentrated *in vacuo*.

Products were characterized by analytical reverse phase HPLC using a Vydac C4 column (5 μ m 300Å, 1.0x150 mm) on a MAGIC 2002 liquid chromatography system (Michrom BioResource, Auburn, CA) and electrospray ionization mass spectrometry (Micromass, FISIONS Instruments, Bevely, MA).

5 Chromatography of the reaction products from part A(2) of this example generated large sharp peaks, which correspond to the lipid-conjugated polyamide compounds, and much smaller peaks, which correspond to unreacted oligomers. Mass spectroscopy confirmed the theoretical molecular weights of the synthesized lipid-conjugated polyamide compounds from part A(2) in this example. Thus, these results
10 confirm the predicted molecular weights of the lipid-conjugated polyamide compounds.

Example 2

Preparation of Lipid-Conjugated Polyamide Compounds Complexed with Plasmid DNA

Lipid-conjugated polyamides (from Example 1) complexed with plasmid DNA
15 were prepared. The plasmid used in these experiments was pCMVkmLUC. Plasmid pCMVkmLUC was constructed by inserting the luc⁺ gene from pSP-luc⁺ (Promega Corp. Madison, WI) into the expression vector pCMVkm2. The sequence of vector CMVkm2 is depicted in SEQ ID NO: 1. A plasmid map of vector CMVkm2 is shown in Figure 2.

20 Briefly, pSP-luc⁺ was digested with the restriction enzymes NheI-EcoRV (Boehringer Mannheim, Indianapolis, IN) and a fragment of 1691 bp was isolated by standard methods. This fragment was inserted into pCMVkm2, which had been digested with XbaI, EcoRV using the Rapid Ligation Kit (Boehringer Mannheim, Indianapolis, IN). The luc⁺ gene was cloned into pCMVkm2 such that expression is driven by the
25 CMV immediate early enhancer promoter and terminated by the bovine growth hormone polyadenylation signal.

Lipid-conjugated polyamide compounds synthesized in Example 1 were dissolved in sterile water at a concentration of 5 mM and sonicated for 1 minute to get a clear solution. A 5 mM solution of pCMVkmLUC in sterile water was prepared. Equal
30 volumes of the lipid-conjugated polyamide compound solution and plasmid solution were mixed together and allowed to sit for 15 minutes at room temperature before

transfection. For transfection studies, OptiMEM (GibcoBRL, Gaithersburg, MD) was used instead of sterile water.

Complexes were also prepared by mixing lipid-conjugated polyamide compounds with DOPE or cholesterol in an equal molar ratio, then sonicated to form liposomes.

5 Plasmid DNA was then added to these preformed liposomes to form complexes.

Complexes of Lipofectin[®] (GibcoBRL, Gaithersburg, MD) and DMRIE-C[™] (GibcoBRL, Gaithersburg, MD) and plasmid DNA were also prepared as controls in the following examples, in accordance with manufacturer's directions.

10

Example 3

Characterization of Lipid-Conjugated Polyamide Compounds Complexed with Plasmid DNA

Rehydration of the lyophilized Compound 16 (see Table 2) in sterile water produced a clear solution which did not scatter light. Under negative-stain electron
15 microscopy, the lipid-conjugated polyamide compound in this solution appeared as aggregates of most cylindrical micelles, with some spherical micelles with diameters of between about 10 to about 15 nm. While mixing a solution of Compound 16 with the pCMVkmLUC solution from Example 3, dynamic light scattering measurements (N4 Plus, Coulter Instruments, Miami, FL) indicated that particle sizes reached a minimum of
20 about 120 nm when the +/- charge ratio was between about 2 and about 4. The particles sizes of these complexes increased slightly as the charge ratio changed. Negative-stain electron microscopy of a 2:1 +/- charge ratio complex indicated the formation of homogeneous, spherical particles with diameters of around 150 nm.

Dynamic light scattering measurements were similarly taken while mixing
25 Compound 23 with pCMVkmLUC. The particle sizes of the resulting lipid-conjugated polyamide/DNA complex reached a minimum of 120 nm when the +/- charge ratio was between about 2 and about 4. When the charge ratio increased or decreased, the particle sizes of these complexes increased slightly.

Zeta potential measurements (Delsa 400, Coulter Instruments) indicated the zeta
30 potential of the Compound 16/DNA complex increased steadily as the ratio of lipid-

conjugated polyamide to DNA increased, and charge neutralization was realized at a charge ratio between about 0.5 and 1.

Complex formation was also characterized by agarose gel mobility shift assay. Complexes of Compound 16 with 1 μ g pCMVkmLUC at a +/- charge ratio of 1 or above, were loaded onto a 1% agarose gel (70V, 1 hour) to examine the retardation of the complexed plasmid DNA, as compared to naked plasmid DNA. Gel mobility shift confirmed that the plasmid DNA was retained in the complex.

Example 4

10 Characterization of DNA Stability in Complexes with Lipid-Conjugated Polyamide Compounds

To determine whether lipid-conjugated polyamide compounds of the present invention inhibit degradation of complexed DNA by DNase I (Boehringer Mannheim, Indianapolis, IN), lipid-conjugated polyamide/DNA complexes were treated with DNase I and the results were analyzed by agarose gel electrophoresis. pCMVkmLUC (10 μ g) and lipid-conjugated/pCMVkmLUC complexes containing 10 μ g of pCMVkmLUC were incubated at 37°C with 10 units of DNase I in 50 μ l of 10 mM MgCl₂ for 15 minutes. An aliquot (1 μ g DNA) of complex/DNase mixture was loaded (the loading buffer
20 contained 1% SDS) onto a 1% agarose gel (70V, 1 hour) to examine the integrity of the plasmid DNA.

The results indicated that lipid-conjugated polyamide compounds having either aromatic or aliphatic R₁ groups were effective at providing resistance to DNase degradation, however, aromatic R₁ groups appeared to provide somewhat greater
25 resistance to degradation. These results suggest that lipid-conjugated polyamide compounds having aromatic R₁ groups complex more strongly with DNA.

Furthermore, lipid-conjugated polyamide compounds of formula (IV), having the R₁¹:R₁²:R₁³ motif of aminoethyl:2-(4'-methoxyphenyl)ethyl:2-(4'-methoxyphenyl)ethyl, appeared to protect the plasmid DNA to a greater extent, than other motifs, as evidenced
30 by the greater amount of supercoiled plasmid DNA retained. This result appeared to be independent of the length of the lipid-conjugated polyamide compound. Plasmid DNA stability was also evaluated as a function of complex +/- charge ratio.

Lipitoid/polynucleotide complexes having different charge ratios were prepared by mixing different amounts of Compound 16 with a fixed amount of plasmid DNA. No significant degradation was observed with any of the charge ratios evaluated, i.e., +/- 10:1, 8:1, 4:1 and 2:1. However, supercoil conformation was maintained to a greater extent at higher charge ratios as compared to lower charge.

Example 6

Transfection Method and Assay

10

Three cell lines were used in the transfection studies, HT1080 (American Type Culture Collection, Rockville, MD, Accession No. CCL 121), NIH3T3 (from culture collection stock of Chiron Corp., Emeryville, CA), and Cos6M (from laboratory of E. Glazer, Chiron Corp. Emeryville, CA). Prior to transfection, the cells (2 ml of a suspension of 1×10^5 cells/ml DME-FCS) were plated into each well of a 6-well plate (Corning, Cambridge, MA) and moved gently to evenly disperse the cells. The cells were transfected 24 hours later.

15

For transfection, both transient and stable complexes containing plasmid (GibcoBRL, Gaithersburg, MD). A transfection medium of complexed DNA was added to each well at 1 μ g pCMVkmLUC/well, and at a concentration of 1 μ g/100 μ l. All tested conditions were performed in duplicate wells. Cells were cultured with transfection medium for 3 hours at 37 °C. The transfection medium was removed by pipet without disturbing the cell layer and replaced with 2 ml of either DME-FCS (for serum-positive conditions) or OptiMEM (for serum-free conditions). The cells were reincubated for 48 hours, then the medium discarded. The cells were then rinsed twice with DPBS (JRH Biosciences, Lenexa, KS)

20

25

Reporter lysis buffer (300 μ l/well) (Promega Corp., Madison, WI) was added to the wells and the cells were allowed to lyse for 15 to 20 minutes on a rocker. The cell lysate from each well was transferred to an eppendorf tube using a cell scraper (Fisher Scientific, Pittsburgh, PA), followed by a freeze (ethanol-dry ice)-thaw cycle and microfugation at maximum speed for 2 minutes.

30

Firefly luciferase activity for the lysates was assayed using the Luciferase Assay Kit (Promega Corp. , Madison, WI) and an ML2250 automated luminometer (Dynatech, Chantilly, VA) in accordance with manufacturer's directions.

5

Example 7

Evaluation of Transfection Efficiency as a Function of Lipid-Conjugated Polyamide/DNA Complex Charge Density

10 Lipid-conjugated polyamide/DNA complexes were prepared from Compound 16 and pCMVkmLUC in OptiMEM as described in Example 2, which had +/- charge densities of 0.5:1, 1:1, 2:1, 4:1, and 8:1. HT 1080 cells were transfected and assayed in accordance with the method described in Example 6. The relationship between transfection efficiency, i.e., luciferase activity, and complex charge density is shown in
15 Figure 2. The results indicate that transfection of HT1080 cells using Compound 16 was most efficient when the +/- charge density of the complex was about 2:1. The results also indicate that serum did not appear to have a significant influence on transfection efficiency.

20

Example 8

Evaluation of the Effect of Oligomer Length on the Transfection Efficiency of Lipid-Conjugated Polyamide Compounds

Lipid-conjugated polyamide/pCMVkmLUC complexes were prepared from
25 Compounds 15-19, having a +/- charge ratio of 2:1, in OptiMEM according to Example 2. Compounds 15-19 all have the same general structure, i.e. formula (IV), identical side chains and lipid groups, however, they differ with respect to oligomer length, i.e., the integer m is different for each of these compounds. The integer m is 2, 3, 4, 8, and 12, for Compounds 15, 16, 17, 18, and 19, respectively. HT1080 cells were transfected with
30 complexes prepared from each of these compounds, then assayed as described in Example 6.

The results, shown in Figure 3, indicate that for this particular series of compounds, transfection efficiency is highest for Compound 16 (m=3).

Example 9

Effect of Lipid Moiety Type on Transfection Efficiency

Conjugated-lipid polyamide/pCMVkmLUC complexes were prepared from
5 Compounds 16 and 23, both DMPE-conjugated compounds, and Compounds 20 and 24,
both DOPE-conjugated compounds, having a +/- charge density ratio of 2:1, as described
in Example 2. HT1080 cells were transfected using these complexes and transfection
efficiency was assayed, both in accordance with the methods described in Example 6

The results, shown in Figure 4, indicated higher transfection efficiencies for the
10 DMPE-conjugated compounds, as compared to the DOPE-conjugated compounds.

Example 10

Transfection of Different Cell Lines Using Lipid-Conjugated Polyamide Compounds

Lipid-conjugated polyamide/pCMVkmLUC plasmid complexes were prepared
15 from Compounds 16 and 23 in OptiMEM, having a +/- charge density ratio of 2:1,
according to Example 2. In addition, Lipofectin[®]/pCMVkmLUC and DMRIE-
C[™]/pCMVkmLUC complexes were prepared according to manufacturer's directions in
OptiMEM. HT1080, Cos6M, and NIH3T3 cells were transfected with the complexes,
and subsequently assayed for luciferase activity according to Example 6.

20 The results are shown in Table 3. These results demonstrate the higher
transfection efficiencies of lipid-conjugated polyamide compounds, as compared to
commercially available transfection preparations.

TABLE 3. Transfection Efficiency[†] of Lipid-Conjugated Polyamide Compounds

5	Cell Line	<u>Transfection Vehicle</u>		
		Lipofectin [®]	DMRIE-C [™]	Compound 16 Compound 23
	HT1080	1/1 ^a	5.2/3 ^a	10/7 ^a 17.6/12.8 ^a
	COS6M	1/1 ^a	8/27 ^a	/26 ^a 3/29 ^a
	NIH3T3		1/1 ^b	100/33 ^b 0.76/4.9 ^b

[†] Normalized Luciferase Activity (optiMEM, serum-free)/Normalize Luciferase Activity (serum)
^a Both serum and serum-free activities are normalized to the corresponding luciferase activities from Lipofectin[®]-mediated transfection.
^b Both serum and serum-free activities are normalized to the corresponding luciferase activities from DMRIE-C[™]-mediated transfection.

15

Example 11Transfection of HT1080 Cells Using Various Lipid-Conjugated Polyamide Compounds

Lipid-conjugated polyamide/pCMVkmLUC plasmid complexes were prepared from some of the lipid-conjugated polyamide compounds from Example 1, in OptiMEM as described in Example 2. The complexes had a +/- charge density ratio of 2:1. In addition, Lipofectin[®]/pCMVkmLUC and DMRIE-C[™]/pCMVkmLUC complexes were prepared according to manufacturer's directions in OptiMEM. HT1080 cells were transfected with the complexes, and subsequently assayed for luciferase activity according to Example 6.

The results are shown in below in Table 4.

TABLE 4. Transfection Efficiency[†] of Lipid-Conjugated Polyamide Compounds

Transfection Vehicle	Transfection Efficiency
Lipofectin [®]	1 / 1
DMRIE-C [™]	5.2 / 3
Compound 2	2 / 0.2
Compound 3	1.6 / 1.3
Compound 8	0.1 / 0.1
Compound 9	0.1 / 0.1
Compound 13	4.3 / 3.2
Compound 14	5.7 / 4
Compound 15	4.1 / 2.4
Compound 16	10 / 7
Compound 17	1.7 / 1.1
Compound 20	4 / 3
Compound 21	6 / 9.3
Compound 22	3.5 / 6.2
Compound 23	17/12
Compound 24	2 / 7.2

[†] Normalized Luciferase Activity (optiMEM, serum-free)/Normalize Luciferase Activity (serum)

5 ^a All activities were normalized to the corresponding serum and serum-free luciferase activities from Lipofectin[®] -mediated transfection.

10 Lipid-conjugated polyamide compounds having repeating n-mer units with both cationic and neutral sidechains (R₁) were generally more effective at mediating transfection as compared to lipid-conjugated polyamide compounds having only cationic sidechains.

Example 12In vivo Transfection Using a Lipid-Conjugated Polyamide Compound/pCMVkmLUC Complex

5

A preparation of lipid-conjugated polyamide/pCMVkmLUC complex was prepared by mixing an 200 μ l of a solution of 60 μ g of pCMVkmLUC in 200 μ l of D5W (5% dextrose) with 200 μ l of 2.9 mg of Compound 16 in 200 μ l of D5W (5% dextrose), such that the complex +/- charge ratio was 8:1. The final volume of the lipid-conjugated polyamide/DNA complex preparation was 400 μ l. A 400 μ l dosage of the preparation was injected into the tail vein of a Balb/C mouse.

The mouse was sacrificed 24 hours post-injection and the lungs, liver, heart, kidney, and spleen were harvested. The harvested organs were placed in 2.0 ml screwcap tubes in which one third of the volume of the tubes was filled with glass beads (BioSpec Products, Inc., Bartlesville, OK) Please provide source location. The tubes were frozen in liquid nitrogen, then stored at -70°C.

To extract luciferase from the harvested organs, a 300 μ l aliquot of 1 X Reporter Lysis Buffer (Promega Corp., Madison, WI) was added to each tube. The contents of the tubes were then homogenized for 1 minute at 4°C. After adding a 200 μ l aliquot of 1 X RLB to each tube, the tube contents were vortexed for 30 minutes in the cold room. The tubes and their contents were then frozen in an ethanol/dry ice bath, then thawed in a water bath at 20°C, for three cycles. The tubes were then centrifuged at 12,500 rpm for 5 minutes in a cold room. After centrifugation, the supernatant was collected by pipet.

The supernatant was assayed for luciferase activity using the Promega Luciferase assay system and the Dynatech ML2250 automated luminometer (Dynatech, Chantilly, VA) in accordance with the manufacturer's directions.

The results, shown in Figure 5, indicate that luciferase is expressed in mouse lung, liver, and, to a lesser extent, spleen tissue. These results demonstrate the efficacy of lipid-conjugated polyamide compounds at mediating *in vivo* transfection.

Example 13Dilution-Concentration Method for Formulating Stable Preparations of Delivery Vehicle/Polynucleotide Complexes

5

I. Charge Calculations

The concentration of negative charges on DNA was calculated as 3.03 nmol of phosphate per 1 µg of DNA based on an average molecular weight of 330 daltons for each nucleotide. The formula weight of each lipid-conjugated polyamide compound was calculated as semi-trifluoroacetate salt (50% of amino groups form salt with TFA), and the concentration of the lipid-conjugated polyamide compound was determined on the basis of the basis of the lyophilized product from Example 1.

II. Complex Formation

All steps were conducted at ambient temperature. Diagnosis grade purified water (DGPW) was used to prepare all stock solutions. Lipid-conjugated polyamide/pCMVkmLUC complexes were prepared having +/- charge ratios of 0.5:1, 1:1, 2:1, 4:1, and 8:1. A dilute solution of 50 µg/ml polynucleotide in was prepared, corresponding to 151 µM of negative charge. A 5mM solutions of lipid-conjugated polyamide Compound 16 was prepared.

A volume of the DNA solution was quickly added to an equal volume of each lipid-conjugated polyamide solution with gentle agitation. It was observed that slow addition of the two solutions tended to result in the formation of larger complexes and occasional precipitates. Better results were achieved when the DNA solution was added to the lipid-conjugated polyamide solution rather than vice-versa.

The formulations were concentrated using commercially available ultrafiltration membrane concentrator devices with appropriate nominal molecular weight cutoff (e.g., 100 kd). Two milliliters of each dilute lipid-conjugated polyamide/pCMVkmLUC complex preparation (in 30% (v/v) ethanol) were placed in a Centricon® -100 (Amicon Inc., Beverly, MA) and centrifuged at 1000 x g for 30 minutes or until the volume of the retentate contained the complex was approximately 50 µl. The filtrate was removed from the bottom receiver. The retentate was diluted with 2 ml of 5% glucose, and

concentrated to 50 μ l again by repeating the above procedure. This operation was repeated again to make a concentrated, stable preparation containing 1 mg/ml of pCMVkmLUC in 5% glucose. The method can be conducted cold (e.g., at 4°C) or at ambient temperatures. Generally, the final concentration of DNA in the retentate can be as high as 1 mg/ml without any visible precipitation.

Zeta potential measurements were taken of each concentrated preparation at day 1, day 2, and day 8 post-formulation. The results, shown in Figures 6 and 7, indicate that the zeta potential of the complexes in the concentrated preparation remained substantially stable over a period of 8 days.

Dynamic light scattering measurements (N4 Plus, Coulter Instruments, Miami, FL) were taken for each concentrated preparation of lipid-conjugated polyamide/pCMVkmLUC complex. The results, shown in Figures 8 and 9, indicate that the particle sizes of the complexes also remained substantially stable over a period of 8 days.

HT1080 cells were transfected with each concentrated preparation of lipid-conjugated/pCMVkmLUC complex at 2- and 18-days post-formulation (Compound 16) and at 4- and 12-days post-formulation, according to the method described in Example 6. A preparation of DMRIE-CTM/pCMVkmLUC complexes was prepared using conventional methods described in Example 2, and HT1080 cells were transfected. Cell confluence was also measured using a microscope and estimating the percentage of live cells.


The results, shown in Figures 10-12 indicated that luciferase activity in cells transfected with lipid-conjugated polyamide/pCMVkmLUC complexes prepared according to the "concentration" formulation method, was higher than for DMRIE-CTM/pCMVkmLUC complexes prepared according to conventional methods. The results indicate that cell confluence is high for the 0.5:1, 1:1, and 2:1 +/- charge ratio complexes and the DMRIE-CTM, and lower for the 4:1 and 8:1 +/- charge ratio complexes. Doubling the amount of transfection preparation increased luciferase expression in complexes having a 2:1 +/- charge ratio, but decreased expression in complexes having a 4:1 +/- charge ratio, as shown in Figure 14. Figure 14 also shows that cell confluence also decreased when the amount of 4:1 +/- charge ratio complex was doubled.

The effect of using the conventional method for preparing delivery vehicle/DNA complexes, as described in Example 2, was compared to use of the concentration method for preparing delivery vehicle/DNA complex. Lipid-conjugated polyamide compounds (16 and 23)/pCMVkmLUC complexes were prepared according to the conventional method described in Example 2. HT1080 cells were transfected within 30 minutes of formulation. The results are designated "Mixing" in Figure 15. Similarly, lipid-conjugated polyamide compound (16 and 23)/pCMVkmLUC complexes were prepared according to the concentration method followed by transfection of HT1080 cells 1 to 7 days post-formulation. The results are designated "Preformed" in Figures 14 and 15. Complexes of Lipofectin and DMRIE-C with pCMVkmLUC were also prepared using the conventional formulation method described in Example 2. The results, shown in Figure 15, indicate that luciferase activity was higher in cells transfected with complexes formed by the concentration method, than in cells transfected with complexes formed by the conventional method. Cell viability was measured by staining with trypan blue.

While the invention has been described in detail with reference to certain preferred embodiments thereof, it will be understood that modifications and variations are within the spirit and scope of that which is described and claimed.

What is claimed:

1. A compound having the formula:



wherein n is an integer selected from 1 to about 48 and m is an integer selected from about 2 to about 48,

wherein R_1 for each monomeric unit, $\begin{array}{c} R_1 \\ | \\ N - W - C \\ || \\ O \end{array}$, and R_a are independently selected from the group consisting of a hydrogen atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group; -SH; an optionally substituted, branched or straight chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and a lipid moiety that is optionally bonded to linker moiety,

wherein R_1 is not a hydrogen atom for at least one monomeric unit,

wherein R_c is selected from a hydrogen atom; a hydroxy group; an amino group; a hydrazine group; a sulfonyl group; -SH; an optionally substituted, branched or straight chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and a lipid moiety that is optionally bonded to a linker moiety,

wherein when R_1 , R_a , or R_c is an aryl or arylalkyl group having fewer than about 5 carbon atoms in a backbone structure, said backbone structure further comprises one or more heteroatoms atoms,

wherein W for each monomeric unit is independently selected from an optionally substituted, branched or straight chain divalent moiety having from 1 to about 50 atoms and optionally, one or more double or triple bonds in a backbone that contains carbon and optionally contains nitrogen, oxygen, sulfur, and phosphorus,

wherein said lipid moiety is a hydrophobic or amphipathic moiety selected from the group consisting of:

(i) optionally substituted aryl or arylalkyl moieties having from about 14 to about 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and

(ii) optionally substituted, branched or straight chain aliphatic moieties having from about 10 to about 50 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic moieties optionally have one or more double or triple bonds, and

wherein at least one of R_a , R_c , W for a single monomeric unit and R_1 for a single monomeric unit comprises a lipid moiety.

2. The compound according to claim 1, wherein R_1 is not a hydrogen atom for at least two monomeric units.

3. The compound according to claim 1, wherein R_1 is not a hydrogen atom for any of the monomeric units.


4. The compound according to claim 1, wherein each R_1 is independently selected from the group consisting of a cationic, an anionic, and a neutral sidechain.

5. The compound according to claim 1, wherein R_1 is selected from moieties found on naturally-occurring amino acids and on non-naturally-occurring amino acids.

6. The compound according to claim 4, wherein said cationic sidechain is selected from the group consisting of aminoalkyl and derivatives thereof; (S)- α -methylethylenediamino and derivatives thereof; trimethylaminoethyl and derivatives thereof; guanidinoalkyl and derivatives thereof; aminobenzyl and derivatives thereof; pyridinium and derivatives thereof; and isoamyl and derivatives thereof.

7. The compound according to claim 4, wherein said neutral sidechain is selected from the group consisting of (S)- α -methylbenzyl, (R)- α -methylbenzyl, and derivatives thereof; benzyl and derivatives thereof; phenethyl and derivatives thereof; naphthylmethyl and derivatives thereof; (S)- α -methylnaphthyl, (R)- α -methylnaphthyl, and derivatives thereof; N-propylpyrrolidinone and derivatives thereof; cyclohexylmethyl and derivatives thereof; furfuryl and derivatives thereof; 3,4,5-trimethoxybenzyl and derivatives thereof; alkoxyalkyl and derivatives thereof; and p-methoxyphenethyl and derivatives thereof.

8. The compound according to claim 4, wherein said anionic side chain is selected from the group consisting of carboxy, carboxy methyl, carboxy ethyl, and derivatives thereof; benzoic acid and derivatives thereof; phosphates and derivatives thereof; and sulfates and derivatives thereof.

9. The compound according to claim 1, wherein -W- is selected from the group consisting of $-\text{CH}_2\text{CH}_2-$, , $-\text{CH}_2\text{CH}_2-\text{O}-$, and $-\text{CH}_2-\text{CH}=\text{CH}-$.

10. The compound according to claim 1, wherein -W- has the formula
 $-\text{CR}_2\text{R}_3-$,

wherein R_2 and R_3 for each monomeric unit are monovalent moieties independently selected from a hydrogen atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group; $-\text{SH}$; an optionally substituted, branched or straight chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that

optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein said aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and a lipid moiety that is optionally bonded to a linker moiety,

wherein when either R_2 and R_3 is an aryl or arylalkyl group having fewer than about 5 carbon atoms in a backbone structure, said backbone structure further comprises one or more heteroatoms.


11. The compound according to claim 10, wherein R_2 and R_3 are both a hydrogen atom.
12. The compound according to claim 1, wherein R_a comprises a lipid moiety.
13. The compound according to claim 1, wherein at least one R_1 comprises a lipid moiety.
14. The compound according to claim 1, wherein R_c comprises a lipid moiety.
15. The compound according to claim 11, wherein R_a comprises a lipid moiety.
16. The compound according to claim 11, wherein at least one R_1 comprises a lipid moiety.
17. The compound according to claim 11, wherein R_c comprises a lipid moiety.

18. The compound according to claim 1, wherein said lipid moiety is naturally or synthetically derived.

19. The compound according to claim 1, wherein said lipid moiety is derived from phosphoglycerides, glycosylglycerides, sphingolipids, and saturated and unsaturated sterols.

20. The compound according to claim 19, wherein said phosphoglyceride is a phosphatidyl ethanolamine.

21. The compound according to claim 1, wherein n is 3 such that said compound has the formula




wherein R_1^1 is a cationic side chain, and R_1^2 and R_1^3 are both neutral side chains.

22. The compound according to claim 21, wherein R_a comprises a lipid moiety.

23. The compound according to claim 22, wherein said lipid moiety is derived from a compound selected from the group consisting of a phosphoglyceride, a glycosylglyceride, a sphingolipid, a saturated sterol and an unsaturated sterol.

24. The compound according to claim 23, wherein said phosphoglyceride is selected from the group consisting of a phosphatidylethanolamine and a phosphatidylpropanolamine.

25. The compound according to claim 24, wherein said R_a has the formula




wherein p is an integer selected from 2 or 3, and each R_4 is independently selected from an alkyl or alkenyl moiety having from about 6 to about 25 carbon atoms in a backbone.

26. A method of synthesizing lipid-conjugated polyamide compounds, said method comprising:

a) contacting

- (1) a lipid reactant, with
- (2) an oligomer reactant, wherein said oligomer reactant has the

general formula:



wherein n is an integer selected from 1 to about 48, and m is an integer from about 2 to about 48,

wherein each T_a and T_c is independently selected from a terminal group and a reactive moiety that is capable of further reaction with said lipid reactant,

wherein R_1 for each monomeric unit, $\begin{array}{c} R_1 \\ | \\ N - W - C \\ || \\ O \end{array}$, in said oligomer reactant is selected from the group consisting of a hydrogen atom; a hydroxy group; an amino group; a carboxyl group; a sulfonyl group, -SH; an optionally substituted, branched or straight chain aliphatic group having from 1 to about 8 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the

aliphatic group optionally has one or more double or triple bonds; an optionally substituted aryl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus; an optionally substituted arylalkyl group having from about 3 to about 12 carbon atoms in a backbone structure that optionally contains nitrogen, oxygen, sulfur, and phosphorus, wherein the alkyl component of said arylalkyl optionally has one or more double or triple bonds; and a reactive moiety that is capable of further reaction with said lipid reactant,

wherein when R_1 , R_a , or R_c is an aryl or arylalkyl group having fewer than about 5 carbon atoms in a backbone structure, said backbone structure further comprises one or more heteroatoms,

wherein R_1 is not a hydrogen atom for at least one monomeric unit,

wherein W for each monomeric unit is selected from an optionally substituted, branched or straight chain divalent moiety having from 1 to about 50 atoms in a backbone that contains carbon, and optionally contains nitrogen, oxygen, sulfur, and phosphorus, and optionally one or more double or triple bonds,

wherein at least one of T_a , T_c , W for a single monomeric unit, and R_1 for a single monomeric unit comprises a reactive moiety that is capable of further reaction with said lipid reactant; then

b) reacting said lipid reactant with said oligomer reactant to conjugate the lipid reactant to the oligomer reactant.

27. A composition comprising the lipid-conjugated polyamide compound of claim 1 and a biologically active agent.

28. A method for inducing the uptake of a biologically active agent by a cell, said method comprising:

providing a composition comprising an effective amount of a biologically active agent and a lipid-conjugated polyamide compound; then
contacting a biological sample with an effective dose of said composition,
wherein said biological sample comprises a cell.

29. The method according to claim 28, wherein said biologically active agent is a polynucleic acid.

30. A method for inducing the uptake of a biologically active agent by a cell *in vivo*, said method comprising:

providing a composition comprising an effective amount of a biologically active agent and the lipid-conjugated polyamide compound of claim 1; then
administering an effective dose of said composition to a subject.

31. The method according to claim 30, wherein said biologically active agent is a polynucleic acid.

32. A method of expressing a gene in a mammal, said method comprising:
administering a polynucleic acid complexed with the lipid-conjugated polyamide compound of claim 1 to a mammal,

wherein said polynucleic acid is capable of functionally expressing said gene in said mammal, and

wherein said complex is effective at transfecting said gene into a cell in said mammal.

33. A method for substantially inhibiting nuclease-induced polynucleotide degradation, said method comprising:

contacting a polynucleotide with a degradation-inhibiting quantity of the lipid-conjugated polyamide compound of claim 1, and

introducing said polynucleotide and said lipid-conjugated polyamide compound into a nuclease-containing environment.

34. A method of making a stable preparation of a polynucleic acid complexed with a delivery vehicle, said method comprising:

- a) providing a polynucleic acid in a first liquid carrier as a dilute polynucleic acid solution that is substantially precipitant-free;
- b) providing a delivery vehicle-forming compound in a second liquid carrier as a delivery vehicle solution that is substantially precipitant-free;
- c) combining said dilute polynucleic acid solution with said delivery vehicle solution to form a dilute preparation of delivery vehicle/polynucleic acid complex; then
- d) reducing the volume of said dilute preparation to form a stable preparation of delivery vehicle/polynucleic acid complex,

wherein the concentration of polynucleic acid in said stable preparation is higher than the concentration of polynucleic acid in said dilute polynucleic acid solution, and

wherein said stable preparation is substantially precipitant-free.

35. The method according to claim 34, wherein said delivery vehicle is a liposome.

36. The method according to claim 34, wherein said delivery vehicle-forming compound is selected from a lipid, a polycationic non-lipid compound, and a lipid-conjugated polyamide compound.

1/15




FIG. 1

2/15




FIG. 2

3/15





FIG. 3

4/15



Lipid-conjugated polyamide compound

FIG. 4

5/15




FIG. 5

6/15




FIG. 6

7/15




FIG. 7

8/15




FIG. 8

9/15




FIG. 9

10/15




FIG. 10

11/15




FIG. II

12/15




FIG. 12

13/ 15




FIG. 13

14/15




FIG. 14

15/15




FIG. 15

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: CHIRON CORPORATION

(ii) TITLE OF INVENTION: Lipid-Conjugated Polyamide Compounds and Related Compositions and Methods Thereof

(iii) NUMBER OF SEQUENCES: 1

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Chiron Corporation
(B) STREET: 4560 Horton Street
(C) CITY: Emeryville
(D) STATE: California
(E) COUNTRY: U.S.A.
(F) ZIP: 94608-2916

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: Unassigned
(B) FILING DATE: Even date Herewith
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Fujita, Sharon M.
(B) REGISTRATION NUMBER: 38,459
(C) REFERENCE/DOCKET NUMBER: 1387.100

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (510) 923-2706
(B) TELEFAX: (510) 655-3542

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4328 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCCGCGGAAT	TTCGACTCTA	GGCCATTGCA	TACGTGTGAT	CTATATCATA	ATATGTACAT	60
TTATATTGGC	TCATGTCCAA	TATGACCGCC	ATGTTGACAT	TGATTATTGA	CTAGTTATTA	120
ATAGTAATCA	ATTACGGGGT	CATTAGTTCA	TAGCCCATAT	ATGGAGTTCC	GCGTTACATA	180
ACTTACGGTA	AATGGCCCCG	CTGGCTGACC	GCCCAACGAC	CCCCGCCCAT	TGACGTCAAT	240
AATGACGTAT	GTTCCCATAG	TAACGCCAAT	AGGGACTTTC	CATTGACGTC	AATGGGTGGA	300
GTATTTACGG	TAAACTGCCC	ACTTGGCAGT	ACATCAAGTG	TATCATATGC	CAAGTCCGCC	360
CCCTATTGAC	GTCAATGACG	GTAATGGCC	CGCCTGGCAT	TATGCCAGT	ACATGACCTT	420
ACGGGACTTT	CCTACTTGGC	AGTACATCTA	CGTATTAGTC	ATCGCTATTA	CCATGGTGAT	480
GCGGTTTTGG	CAGTACACCA	ATGGGCGTGG	ATAGCGGTTT	GACTCACGGG	GATTTCCAAG	540
TCTCCACCCG	ATTGACGTCA	ATGGGAGTTT	GTTTTGGCAC	CAAAATCAAC	GGGACTTTCC	600
AAAAATGTCGT	AATAACCCCG	CCCCGTTGAC	GCAAAATGGGC	GGTAGGCGTG	TACGGTGGGA	660
GGTCTATATA	AGCAGAGCTC	GTTTAGTGAA	CCGTCAGATC	GCCTGGAGAC	GCCATCCAAG	720
CTGTTTTGAC	CTCCATAGAA	GACACCGGA	CCGATCCAGC	CTCCGCGGCC	GGGAACGGTG	780
CATTGGAACG	CGGATTCCCC	GTGCCAAGAG	TGACGTAAGT	ACCGCCTATA	GACTCTATAG	840
GCACACCCCT	TTGGCTCTTA	TGCATGCTAT	ACTGTTTTTG	GCTTGGGGCC	TATACACCCC	900
CGCTCCTTAT	GCTATAGGTG	ATGGTATAGC	TTAGCCTATA	GGTGTGGGTT	ATTGACCATT	960
ATTGACCACT	CCCCTATTGG	TGACGATACT	TTCCATTACT	AATCCATAAC	ATGGCTCTTT	1020
GCCCAACTA	TCTCTATTGG	CTATATGCCA	ATACTCTGTC	CTTCAGAGAC	TGACACGGAC	1080
TCTGTATTTT	TACAGGATGG	GGTCCATTTA	TTATTTACAA	ATTACATAT	ACAACAACGC	1140
CGTCCCCCGT	GCCCCGAGTT	TTTATTAAAC	ATAGCGTGGG	ATCTCCGACA	TCTCGGGTAC	1200
GTGTTCCGGA	CATGGGCTCT	TCTCCGGTAG	CGGCGGAGCT	TCCACATCCG	AGCCCTGGTC	1260
CCATCCGTCC	AGCGGCTCAT	GGTCGCTCGG	CAGCTCCTTG	CTCCTAACAG	TGGAGGCCAG	1320
ACTTAGGCAC	AGCACAATGC	CCACCACCAC	CAGTGTGCCG	CACAAGGCCG	TGGCGGTAGG	1380
GTATGTGTCT	GAAAATGAGC	TGGGAGATTG	GGCTCGCACC	TGGACGCAGA	TGGAAGACTT	1440
AAGGCAGCGG	CAGAAGAAGA	TGCAGGCAGC	TGAGTTGTTG	TATTCTGATA	AGAGTCAGAG	1500
GTAATCCCG	TTGCCGTGCT	GTTAACGGTG	GAGGGCAGTG	TAGTCTGAGC	AGTACTCGTT	1560
GCTGCCGCGC	GCGCCACCAG	ACATAATAGC	TGACAGACTA	ACAGACTGTT	CCTTTCCATG	1620
GGTCTTTTTCT	GCAGTCACCG	TCGTCGACCT	AAGAATTCAG	ACTCGAGCAA	GTCTAGAAAG	1680
CCATGATATC	CGGATCCACT	ACGCGTTAGA	GCTCGCTGAT	CAGCCTCGAC	TGTGCTTCT	1740
AGTTGCCAGC	CATCTGTTGT	TTGCCCTCC	CCCGTGCCTT	CCTTGACCCT	GGAAGGTGCC	1800
ACTCCCACTG	TCCTTTCCTA	ATAAAATGAG	GAAATTCAT	CGCATTGTCT	GAGTAGGTGT	1860
CATTCTATTG	TGGGGGGTGG	GGTGGGGCAG	GACAGCAAGG	GGGAGGATTG	GGAAGACAAT	1920
AGCAGGGGGG	TGGGCGAAGA	ACTCCAGCAT	GAGATCCCCG	CGCTGGAGGA	TCATCCAGCC	1980
GGCGTCCCGG	AAAACGATTC	CGAAGCCCAA	CCTTTCATAG	AAGGCGGCGG	TGGAATCGAA	2040
ATCTCGTGAT	GGCAGGTTGG	GCGTCGCTTG	GTCGGTCATT	TCGAACCCCA	GAGTCCCGCT	2100
CAGAAGAACT	CGTCAAGAAG	GCGATAGAAG	GCGATGCGCT	GCGAATCGGG	AGCGGCGATA	2160
CCGTAAAGCA	CGAGGAAGCG	GTCAGCCCAT	TCGCCGCCAA	GCTCTTCAGC	AATATCACGG	2220
GTAGCCAACG	CTATGTCCTG	ATAGCGGTCC	GCCACACCCA	GCCGCGCACA	GTCGATGAAT	2280
CCAGAAAAGC	GGCCATTTTC	CACCATGATA	TTCCGGCAAGC	AGGCATCGCC	ATGGGTACAG	2340
ACGAGATCCT	CGCCGTCGGG	CATGCGGCC	TTGAGCCTGG	CGAACAGTTC	GGCTGGCGCG	2400
AGCCCTGAT	GCTCTTCGTC	CAGATCATCC	TGATCGACAA	GACCGGCTTC	CATCCGAGTA	2460
CGTGCTCGCT	CGATGCGATG	TTTCGCTTGG	TGGTCGAATG	GGCAGGTAGC	CGGATCAAGC	2520
GTATGCAGCC	GCCGATTGCG	ATCAGCCATG	ATGGATACTT	TCTCGGCAGG	AGCAAGGTGA	2580
GATGACAGGA	GATCTTGCCC	CGGCACCTCG	CCCAATAGCA	GCCAGTCCCT	TCCCGCTTCA	2640
GTGACAACGT	CGAGCACAGC	TGCGCAAGGA	ACGCCGTCG	TGGCCAGCCA	CGATAGCCCG	2700
GCTGCCTCGT	CCTGCAGTTC	ATTCAGGGCA	CCGGACAGGT	CGGTCTTGAC	AAAAAGAACC	2760
GGGCGCCCT	GCGCTGACAG	CCGGAACACG	GCGGCATCAG	AGCAGCCGAT	TGTCTGTTGT	2820
GCCCAGTCAT	AGCCGAATAG	CCTCTCCACC	CAAGCGGCCG	GAGAACCTGC	GTGCAATCCA	2880
TCCTGTTCAT	TCATGCGAAA	CGATCCTCAT	CCTGTCTCTT	GATCAGATCT	TGATCCCTCG	2940
CGCCATCAGA	TCCTTGGCGG	CAAGAAAGCC	ATCCAGTTTA	CTTTGCAGGG	CTTCCCAACC	3000
TTACCAGAGG	GCGCCCCAGC	TGGCAATTCC	GGTTCGCTTG	CTGTCCATAA	AACCGCCAG	3060

TCTAGCTATC	GCCATGTAAG	CCCACTGCAA	GCTACCTGCT	TTCTCTTTGC	GCTTGCGTTT	3120
TCCCTTGTC	AGATAGCCCA	GTAGCTGACA	TTCATCCGGG	GTCAGCACCG	TTTCTGCGGA	3180
CTGGCTTTCT	ACGTGTTCCG	CTTCCTTTAG	CAGCCCTTGC	GCCCTGAGTG	CTTGCGGCAG	3240
CGTGAAGCTG	TCAATCCGC	GTTAAATTTT	TGTTAAATCA	GCTCATTTTT	TAACCAATAG	3300
GCCGAAATCG	GCAAAATCCC	TTATAAATCA	AAAGAATAGC	CCGAGATAGG	GTTGAGTGTT	3360
GTTCCAGTTT	GGAACAAGAG	TCCACTATTA	AAGAACGTGG	ACTCCAACGT	CAAAGGGCGA	3420
AAAACCGTCT	ATCAGGGCGA	TGGCGGATCA	GCTTATGCGG	TGTGAAATAC	CGCACAGATG	3480
CGTAAGGAGA	AAATACCGCA	TCAGGCGCTC	TTCCGCTTCC	TCGCTCACTG	ACTCGCTGCG	3540
CTCGGTCTGT	CGGCTGCGGC	GAGCGGTATC	AGCTCACTCA	AAGGCGGTAA	TACGGTTATC	3600
CACAGAATCA	GGGGATAACG	CAGGAAAGAA	CATGTGAGCA	AAAGGCCAGC	AAAAGGCCAG	3660
GAACCGTAAA	AAGGCCGCGT	TGCTGGCGTT	TTTCCATAGG	CTCCGCCCCC	CTGACGAGCA	3720
TCACAAAAAT	CGACGCTCAA	GTCAGAGGTG	GCGAAACCCG	ACAGGACTAT	AAAGATACCA	3780
GGCGTTTCCC	CCTGGAAGCT	CCCTCGTGCG	CTCTCCTGTT	CCGACCCTGC	CGCTTACCGG	3840
ATACCTGTCC	GCCTTTCTCC	CTTCGGGAAG	CGTGGCGCTT	TCTCATAGCT	CACGCTGTAG	3900
GTATCTCAGT	TCGGTGTAGG	TCGTTGCTC	CAAGCTGGGC	TGTGTGCACG	AACCCCCCGT	3960
TCAGCCCGAC	CGCTGCGCCT	TATCCGGTAA	CTATCGTCTT	GAGTCCAACC	CGGTAAGACA	4020
CGACTTATCG	CCACTGGCAG	CAGCCACTGG	TAACAGGATT	AGCAGAGCGA	GGTATGTAGG	4080
CGGTGCTACA	GAGTTCTTGA	AGTGGTGGCC	TAACAGGATT	AGCAGAGCGA	GGTATGTAGG	4140
TGGTATCTGC	GCTCTGCTGA	AGCCAGTTAC	CTTCGGAAAA	AGAGTTGGTA	GCTCTTGATC	4200
CGGCAACAA	ACCACCGCTG	GTAGCGGCGG	TTTTTTGTTT	GCAAGCAGCA	GATTACGCGC	4260
AGAAAAAAG	GATCTCAAGA	AGATCCTTTG	ATCTTTTCTA	CTGAACGGTG	ATCCCCACCG	4320
GAATTGCG						4328

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/16970

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K47/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 04001 A (MOLECULAR STRUCTURAL BIOTECHNO) 15 February 1996 see claims 8,33; figure 1; examples 4-8 ---	1-5, 10-20, 27-36
X,P	WO 98 27209 A (UNIV EMORY) 25 June 1998 see claims 54-80; figure 2 ---	1-36
X	WO 94 06451 A (CHIRON CORP) 31 March 1994 cited in the application see page 34, line 10-20	1-36
Y	see page 18; claims 12-22,26,29; example 9; tables 1,5 see page 25, line 22-33 ---	1-36
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

S" document member of the same patent family

Date of the actual completion of the international search

7 January 1999

Date of mailing of the international search report

18/01/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gonzalez Ramon, N

INTERNATIONAL SEARCH REPORT

Int. l. Application No.

PCT/US 98/16970

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	US 5 679 559 A (KIM JIN-SEOK ET AL) 21 October 1997 see column 2, line 1-20; figure 1 see column 3, line 55-68; claims 1-3,6-8,17; examples 1,3,5,8 ---	1-5, 10-20, 27-36
X,P	WO 98 06437 A (CHIRON CORP) 19 February 1998 see the whole document ---	1-36
Y,P	WO 97 45442 A (IMPERIAL COLLEGE ;MILLER ANDREW DAVID (GB); COOPER ROBERT GEOFFREY) 4 December 1997 see abstract; figures 2,22 ---	1-36
X	EP 0 545 016 A (BOEHRINGER INGELHEIM INT ;GENENTECH INC (US); UNIV NORTH CAROLINA) 9 June 1993 see page 17, line 35-55; claims 3,31,33,49,50,55; figure 6; examples 6,13,34 ---	1,4,5, 10-29
X	MERWIN J R ET AL: "TARGETED DELIVERY OF DNA USING YEE(GALNACAH)3, A SYNTHETIC GLYCOPEPTIDE LIGAND FOR THE ASIALOGLYCOPROTEIN RECEPTOR" BIOCONJUGATE CHEMISTRY, vol. 5, no. 6, 1 November 1994, pages 612-620, XP000484176 see page 612, column 2, paragraph 2; figure 2 see page 616, column 2 ---	1-5, 10-20, 27-36
Y,P	ABERLE, A. M. ET AL: "A novel tetraester construct that reduces cationic lipid-associated cytotoxicity. Implications for the onset of cytotoxicity" BIOCHEMISTRY,1998, pages 6533-6540, XP002088531 see discussion see abstract; figures 1,2 ---	1-36
X	ZHOU X ET AL: "DNA transfection mediated by cationic liposomes containing lipopolylysine: characterization and mechanism of action." BIOCHIM BIOPHYS ACTA, JAN 19 1994, 1189 (2) P195-203, XP002088532 NETHERLANDS see the whole document ---	1-5, 10-20, 27-36

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/16970

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>LEGENDRE J Y ET AL: "DIOLEOYLMELITIN AS A NOVEL SERUM-INSENSITIVE REAGENT FOR EFFICIENT TRANSFECTION OF MAMMALIAN CELLS" BIOCONJUGATE CHEMISTRY, vol. 8, no. 1, January 1997, pages 57-63, XP000642730 see abstract; figure 1</p>	1-36
Y,P	<p>CROOK K. ET AL: "Inclusion of cholesterol in DOTAP transfection complexes increases the delivery of DNA to cells in vitro in the presence of serum" GENE THERAPY, vol. 5, 1998, pages 137-143, XP002089306 see abstract</p>	1-36

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/16970

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9604001 A	15-02-1996	US 5716614 A AU 3275595 A	10-02-1998 04-03-1996
WO 9827209 A	25-06-1998	AU 5729898 A	15-07-1998
WO 9406451 A	31-03-1994	AU 679945 B AU 5292093 A BG 99521 A CA 2144067 A EP 0671928 A FI 951356 A HU 72614 A JP 8501565 T NO 950682 A NZ 256952 A PL 308204 A US 5831005 A	17-07-1997 12-04-1994 31-01-1996 31-03-1994 20-09-1995 26-04-1995 28-05-1996 20-02-1996 18-04-1995 27-07-1997 24-07-1995 03-11-1998
US 5679559 A	21-10-1997	AU 3802397 A WO 9800556 A	21-01-1998 08-01-1998
WO 9806437 A	19-02-1998	NONE	
WO 9745442 A	04-12-1997	AU 2911197 A	05-01-1998
EP 0545016 A	09-06-1993	AU 671084 B AU 2652692 A BG 98718 A BR 9206559 A CA 2118816 A CZ 9400746 A WO 9307283 A EP 0607206 A FI 941474 A HU 71312 A HU 9500694 A JP 10506001 T MX 9205543 A NO 941154 A NZ 244306 A SG 44680 A SK 36894 A US 5547932 A ZA 9207460 A CN 1070946 A	15-08-1996 03-05-1993 28-02-1995 08-11-1994 31-03-1993 17-05-1995 15-04-1993 27-07-1994 30-03-1994 28-11-1995 29-01-1996 16-06-1998 01-05-1993 29-03-1994 26-07-1995 19-12-1997 10-08-1994 20-08-1996 21-02-1994 14-04-1993

